# **Product Datasheet**

# IFIT2 Antibody - BSA Free NBP1-31164

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-31164

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-31164



# NBP1-31164

IFIT2 Antibody - BSA Free

IFT12 Antibody - BSA Free	
Product Information	
Unit Size	100 ul
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Thimerosal
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	0.1M Tris-Glycine (pH7), 10% Glycerol
Target Molecular Weight	55 kDa
Product Description	
Description	Novus Biologicals Rabbit IFIT2 Antibody - BSA Free (NBP1-31164) is a polyclonal antibody validated for use in WB and ICC/IF. Anti-IFIT2 Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	3433
Gene Symbol	IFIT2
Species	Human
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of human IFIT2. The exact sequence is proprietary.
<b>Product Application Details</b>	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:500-1:3000, Immunocytochemistry/ Immunofluorescence 1:100-


<b>Product Application Details</b>	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:500-1:3000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000




#### **Images**

Western Blot: IFIT2 Antibody [NBP1-31164] - p115 and Sec16A are required for optimal IRF-3 activation in response to activation of cytosolic RNA and DNA sensors.HeLa cells were infected with lentiviral vectors encoding shRNA targeting p115. Image collected and cropped by CiteAb from the following publication


(//dx.plos.org/10.1371/journal.ppat.1002747) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: IFIT2 Antibody [NBP1-31164] - HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: IFIT2 protein stained by IFIT2 antibody [N1], N-term diluted at 1:200. Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody diluted at 1:10000. Blue: Hoechst 33342 staining.



Western Blot: IFIT2 Antibody [NBP1-31164] - Sample(30 ug of whole cell lysate)A:A431 B:Hep G2 7.5% SDS PAGE, antibody diluted at 1:1000.



Western Blot: IFIT2 Antibody [NBP1-31164] - p115 & Sec16A are required for optimal IRF-3 activation in response to activation of cytosolic RNA & DNA sensors.HeLa cells were infected with lentiviral vectors encoding shRNA targeting p115 (A) or Sec16A (B) & non-targeting (NT) control shRNA for 24 h followed by puromycin selection (1.5 µg/ml) for 5 days. Cells were left untreated or stimulated with poly I:C (1 µg/ml), poly dA:dT (1 µg/ml) or SeV (200 HAU/ml) for 16 h. Whole-cell lysates were prepared & subjected to immunoblot analysis with indicated antibodies. One of two independent experiments with similar results is shown. Image collected & cropped by CiteAb from the following publication (https://pubmed.ncbi.nlm.nih.gov/22792062), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



#### **Publications**

Rex EA, Seo D, Chappidi S, Pinkham C et Al. FEAR antiviral response pathway is independent of interferons and countered by poxvirus proteins Nat Microbiol 2024-03-28 [PMID: 38538832]

Khowal S, Monga S, Naqvi S et al. Molecular winnowing, expressional analyses and interactome scrutiny of cellular proteomes of oral squamous cell carcinoma Advances in Cancer Biology - Metastasis 2021-10-01

Khan KA TRK-Fused Gene (TFG), a protein involved in protein secretion pathways, is an essential component of the antiviral innate immune response PloS Pathog 2021-01-07 [PMID: 33411856] (WB, WB, Human)

Khan KA, Do F, Marineau A et al. Fine-Tuning of the RIG-I-Like Receptor/Interferon Regulatory Factor 3-Dependent Antiviral Innate Immune Response by the Glycogen Synthase Kinase 3/beta-Catenin Pathway. Mol Cell Biol 2015-09-01 [PMID: 26100021]

Baril M, Es-Saad S, Chatel-Chaix L et al. Genome-wide RNAi Screen Reveals a New Role of a WNT/CTNNB1 Signaling Pathway as Negative Regulator of Virus-induced Innate Immune Responses. PLoS Pathog 2013-06-01 [PMID: 23785285] (WB, Human)

van Zuylen WJ, Doyon P, Clement JF et al. Proteomic profiling of the TRAF3 interactome network reveals a new role for the ER-to-Golgi transport compartments in innate immunity. PLoS Pathog 2012-01-01 [PMID: 22792062] (WB, Human)





## **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP1-31164**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00003433-P01-10ug Recombinant Human IFIT2 GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-31164

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

