

Product Datasheet

HDAC3 Antibody - BSA Free

NBP1-31196

Unit Size: 100 ul

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-31196

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-31196



NBP1-31196

HDAC3 Antibody - BSA Free


Product Information	
Unit Size	100 ul
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 20% Glycerol
Target Molecular Weight	49 kDa

Product Description	
Host	Rabbit
Gene ID	8841
Gene Symbol	HDAC3
Species	Human, Mouse, Rat, Drosophila
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human HDAC3. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunoprecipitation 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:1000, Chromatin Immunoprecipitation (ChIP) Assay dependent

Images


Immunocytochemistry/Immunofluorescence: HDAC3 Antibody [NBP1-31196] - SK-N-SH cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: HDAC3 protein stained by HDAC3 antibody [C3], C-term diluted at 1:400. Red: Phalloidin, a cytoskeleton marker, diluted at 1:200. Scale bar = 10 um.




Immunohistochemistry-Paraffin: HDAC3 Antibody [NBP1-31196] - SW480 xenograft, using HDAC3 antibody at 1:500 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.




Chromatin Immunoprecipitation: HDAC3 Antibody [NBP1-31196] - Sample: 293T whole cell lysate/extract A. 5 ug preimmune rabbit IgG, B. 5 ug of HDAC3 antibody The precipitated DNA was detected by PCR with primer set targeting to p21 promoter.




Western Blot: HDAC3 Antibody [NBP1-31196] - Analysis of on: A: 293T B: A431 C: H1299 D: HeLa S3 E: Hep G2 F: MOLT4 G: Raji 10% SDS PAGE diluted at 1:1000




Western Blot: HDAC3 Antibody [NBP1-31196] - Sample: 1000 ug 293T whole cell lysate/extract A. 50 ug 293T whole cell lysate/extract, B. Control with 2 ug of preimmune rabbit IgG, C. Immunoprecipitation of HDAC3 protein by 2 ug of HDAC3 antibody 10% SDS-PAGE gel.




Western Blot: HDAC3 Antibody [NBP1-31196] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with HDAC3 antibody [C3], C-term. An HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Western Blot: HDAC3 Antibody [NBP1-31196] - Various whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with HDAC3 antibody [C3], C-term diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: HDAC3 Antibody [NBP1-31196] - HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: HDAC3 protein stained by HDAC3 antibody [C3], C-term diluted at 1:500. Blue: Hoechst 33342 staining.




Western Blot: HDAC3 Antibody [NBP1-31196] -Whole cell extract (30 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with HDAC3 antibody [C3], C-term diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Western Blot: HDAC3 Antibody [NBP1-31196] - Whole cell extract (30 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with HDAC3 antibody [C3], C-term diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Western Blot: HDAC3 Antibody [NBP1-31196] - Various whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with HDAC3 antibody [C3], C-term (NBP1-31196) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-31196

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00008841-P01-10ug	Recombinant Human HDAC3 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-31196

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

