

Product Datasheet

BRIP1/FANCI Antibody - BSA Free NBP1-31883

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 8

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-31883

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-31883



NBP1-31883

BRIP1/FANCJ Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Thimerosal
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 20% Glycerol
Target Molecular Weight	141 kDa

Product Description	
Host	Rabbit
Gene ID	83990
Gene Symbol	BRIP1
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 26490168).
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human BRIP1/FANCJ. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Proximity Ligation Assay
Recommended Dilutions	Western Blot 1:500-1:3000, Simple Western, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Proximity Ligation Assay Reported in scientific literature (PMID:33445471)

Images


Western Blot: BRIP1/FANCJ Antibody [NBP1-31883] - Whole cell extract (30 ug) was separated by 5% SDS-PAGE, and the membrane was blotted with FANCJ antibody [N1N2], N-term diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.




Immunocytochemistry/Immunofluorescence: BRIP1/FANCJ Antibody [NBP1-31883] - 293T cells were fixed in 2% paraformaldehyde/culture medium at 37 degrees C for 30 min. Green: BRIP1 protein stained by FANCJ antibody [N1N2], N-term diluted at 1:500. Blue: Hoechst 33343 staining.




Immunohistochemistry-Paraffin: BRIP1/FANCJ Antibody [NBP1-31883] - SAS xenograft. FANCJ antibody [N1N2], N-term dilution: 1:500. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.




Immunocytochemistry/Immunofluorescence: BRIP1/FANCJ Antibody [NBP1-31883] - Expression of BRIP1/FANCJ at the mRNA (graphs) and protein (photos) levels in iPSCs. Gene expression was assessed with RNA-seq in FACS-sorted bone marrow HSCs or with real-time RT-PCR in cultured iPSCs. The box-and-whiskers graphs show the median, 25th and 75th percentile (box) minimum, and maximum values (whiskers) of the FPKM (fragments per kilobase million) or relative expression in comparison to house-keeping genes (geometric mean for HPRT, B2M, and beta-actin), $n = 4$, * $p < 0.05$, ** $p < 0.01$, Mann-Whitney test (HSCs) or Wilcoxon test (iPSCs). Nuclei are labeled with DAPI. Scale bars: 20 μm . Image collected and cropped by CiteAb from the following publication ([//pubmed.ncbi.nlm.nih.gov/33445471/](https://pubmed.ncbi.nlm.nih.gov/33445471/)) licensed under a CC-BY license.




Immunocytochemistry/ Immunofluorescence: BRIP1/FANCJ Antibody [NBP1-31883] - In situ proximity ligation assay (PLA) in HEK293T cells. (A) G4-BRIP1 & (B) G4-HMOX1 interactions (red dots) were visualized in the control group & after DNase or RNase digestion. Most of the G4 interactions were localized in the nucleus (arrows), but some were also present in the cytoplasm. Cells that were fixed in methanol & mouse 1H6 antibodies were used. Nuclei were counterstained with DAPI (blue), where the transmission bright-field images (BF) show the cell morphologies. Scale bar: 10 μm . Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33445471/>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: BRIP1/FANCJ Antibody [NBP1-31883] - Detection of the PLA signal in sorted hematopoietic stem cells. (A) G4-Brip1 & (B) G4-Hmox1 interactions (green dots) were visualized in HSCs & GMP & MPP cells. G4 interactions were visible in the nucleus (arrows) & in the cytoplasm. Nuclei were counterstained with DAPI (blue), where the transmission bright-field images (BF) show the cell morphologies. Cells were fixed in PFA & goat 1H6 antibodies were used. Scale bars: 5 μ m. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33445471>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: BRIP1/FANCJ Antibody [NBP1-31883] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with BRIP1/FANCJ antibody [N1N2], N-term (NBP1-31883) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Publications

Boavida A, Napolitano LM, Santos D et al. FANCJ DNA helicase is recruited to the replisome by AND-1 to ensure genome stability EMBO Rep 2024-01-02 [PMID: 38177925]

Tom Egger, Laura Morano, Marie-Pierre Blanchard, Jihane Basbous, Angelos Constantinou Spatial organization and functions of Chk1 activation by TopBP1 biomolecular condensates. Cell reports 2024-04-04 [PMID: 38578830]

Esin Isik, Kaustubh Shukla, Michaela Pospisilova, Christiane König, Martin Andrs, Satyajeet Rao, Vinicio Rosano, Jana Dobrovolna, Lumir Krejci, Pavel Janscak Mut β -Mut β -FANCJ axis mediates the restart of DNA replication after fork stalling at cotranscriptional G4/R-loops. Science advances 2024-02-14 [PMID: 38324687]

Yaneva D, Sparks JL, Donsbach M et al. The FANCJ helicase unfolds DNA-protein crosslinks to promote their repair Molecular cell 2023-01-05 [PMID: 36608669] (WB, Human)

Summers PA, Lewis B, Gonzalez-Garcia J, Lim AHM Visualising G-quadruplex DNA dynamics in live cells by fluorescence lifetime imaging microscopy Nat Commun 2021-01-09 [PMID: 33420085]

Krzepkowski W, Chudy P, Sokołowski G, et al. Proximity Ligation Assay Detection of Protein-DNA Interactions-Is There a Link between Heme Oxygenase-1 and G-quadruplexes? Antioxidants (Basel, Switzerland) 2021-01-12 [PMID: 33445471] (PLA, Mouse)

kim D, Liu Y, Oberly S et al. ATR-mediated proteome remodeling is a major determinant of homologous recombination capacity in cancer cells. Nucleic Acids Res. 2018-07-12 [PMID: 30010936] (WB, Human)

Sun X, Brieno-Enriquez MA, Cornelius A et al. FancJ (Brip1) loss-of-function allele results in spermatogonial cell depletion during embryogenesis and altered processing of crossover sites during meiotic prophase I in mice. Chromosoma. 2015-10-21 [PMID: 26490168] (WB, Mouse)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-31883

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-14361PEP	BRIP1/FANCJ Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-31883

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

