# **Product Datasheet**

# PCTAIRE2 Antibody - BSA Free NBP1-42688

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-42688

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-42688



## NBP1-42688


PCTAIRE2 Antibody - BSA Free

| Unit Size 100 ul  Concentration 1.0 mg/ml  Storage Store at 4C. Do not freeze.  Clonality Polyclonal  Preservative 0.09% Sodium Azide  Isotype IgG  Purity Immunogen affinity purified  Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description  Host Rabbit  Gene ID 5128  Gene Symbol CDK17  Species Human  Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin (Negative)  | PCTAIREZ ATIIDOUY - BSA FTEE |   |
|--|------------------------------|---|
| Concentration 1.0 mg/ml Storage Store at 4C. Do not freeze.  Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description Host Rabbit Gene ID 5128 Gene Symbol CDK17 Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative) Recommended Dilutions Western Blot 1:2000-1:2000, Immunohistochemistry 1:500-1:2000, Immunoprecipitation (Negative) Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue | Product Information          |   |
| Storage Store at 4C. Do not freeze.  Clonality Polyclonal  Preservative 0.09% Sodium Azide  Isotype IgG  Purity Immunogen affinity purified  Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description  Host Rabbit  Gene ID 5128  Gene Symbol CDK17  Species Human  Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue            | Unit Size                    | 100 ul  |
| Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description Host Rabbit Gene ID 5128 Gene Symbol CDK17 Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue              | Concentration                | 1.0 mg/ml   |
| Preservative 0.09% Sodium Azide  Isotype IgG  Purity Immunogen affinity purified  Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description  Host Rabbit  Gene ID 5128  Gene Symbol CDK17  Species Human  Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue                           | Storage                      | Store at 4C. Do not freeze.                                       |
| Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description Host Rabbit Gene ID 5128 Gene Symbol CDK17 Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative) Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Clonality                    | Polyclonal  |
| Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description  Host Rabbit Gene ID 5128 Gene Symbol CDK17  Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Preservative                 | 0.09% Sodium Azide  |
| Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0)  Product Description  Host Rabbit  Gene ID 5128  Gene Symbol CDK17  Species Human  Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Isotype                      | IgG   |
| Product Description  Host Rabbit  Gene ID 5128  Gene Symbol CDK17  Species Human  Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Purity                       | Immunogen affinity purified                                       |
| Host Gene ID 5128  Gene Symbol CDK17  Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Buffer                       | Tris-Citrate/Phosphate (pH 7.0 - 8.0)                             |
| Gene Symbol CDK17  Species Human Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative) Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Product Description          |   |
| Gene Symbol CDK17  Species Human The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Host                         | Rabbit  |
| Human  The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications  Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions  Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes  Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Gene ID                      | 5128  |
| Immunogen The immunogen for this product maps to a region between residue 1 and 50 of human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Gene Symbol                  | CDK17   |
| human PCTAIRE protein kinase 2 using the numbering given in entry NP_002586.2 (GeneID 5128).  Product Application Details  Applications  Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions  Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes  Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Species                      | Human   |
| Applications  Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation (Negative)  Recommended Dilutions  Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes  Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue   | Immunogen                    | human PCTAIRE protein kinase 2 using the numbering given in entry |
| Immunoprecipitation (Negative)  Recommended Dilutions  Western Blot 1:2000-1:10000, Immunohistochemistry 1:500-1:2000, Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Application Notes  Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Product Application Details  |   |
| Application Notes Immunohistochemistry-Paraffin 1:500-1:2000, Immunoprecipitation (Negative)  Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue  | Applications                 |   |
|  | Recommended Dilutions        |   |
|  | Application Notes            |   |




### **Images**

Western Blot: PCTAIRE2 Antibody [NBP1-42688] - Whole cell lysate from Jurkat (5, 15 and 50 ug) and Ramos (R; 50 ug) cells. Antibody used at 0.4 ug/ml.



Immunohistochemistry-Paraffin: PCTAIRE2 Antibody [NBP1-42688] - Sample: FFPE section of human breast carcinoma (left) and kidney clear cell carcinoma (right). Antibody: Affinity purified rabbit anti- PCTAIRE 2 used at a dilution of 1:1,000 (1ug/ml). Detection: DAB





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP1-42688**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-87316PEP PCTAIRE2 Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-42688

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

