## **Product Datasheet**

# Prokineticin R2/PROKR2 Antibody NBP1-45218

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-45218

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-45218



### NBP1-45218


Prokineticin R2/PROKR2 Antibody

FTOKITIETICIT NZ/FNORNZ AHUBOUY	
Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Product Description	
Host	Goat
Gene ID	128674
Gene Symbol	PROKR2
Species	Human, Rat
Immunogen	Peptide with sequence C-NGNTSFTPNFN corresponding to internal region according to NP_658986.1.
Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA
Recommended Dilutions	Western Blot 0.3-1 ug/ml, Immunohistochemistry 3.75 ug/ml, Immunohistochemistry-Paraffin 3.75 ug/ml, Peptide ELISA Detection limit 1:32000
Application Notes	Western blot: Approx 45kDa band observed in Human Brain (Cerebellum), Testis and Thyroid Gland and in Rat Brain, Stomach and Testis lysates (calculated MW of 44.0kDa according to Human NP_658986.1 and 47.9kDa according to Rat NP_620434.1). Primary incubation was 1 hour IHC: Paraffin embedded Human Brain (Cortex).






Western Blot: Prokineticin R2/PROKR2 Antibody [NBP1-45218] - 0.1ug/ml staining of Human Cer(A), Rat Brain (B) and Rat Testis (C) lysate (35ug protein in RIPA buffer). Detected by chemiluminescence.



Immunohistochemistry-Paraffin: Prokineticin R2/PROKR2 Antibody [NBP1-45218] - 3.75ug/ml staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Western Blot: Prokineticin R2/PROKR2 Antibody [NBP1-45218] - (1ug/ml) staining of Rat Stomach lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



#### **Publications**

Monnier C, Dode C, Fabre L et al. PROKR2 missense mutations associated with Kallmann syndrome impair receptor signalling activity. Hum Mol Genet 2009-01-01 [PMID: 18826963]





#### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### **Products Related to NBP1-45218**

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

NBP1-92290PEP Prokineticin R2/PROKR2 Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-45218

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



