

Product Datasheet

STAP2 Antibody NBP1-51904

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-51904

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-51904



NBP1-51904**STAP2 Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat STAP2 Antibody (NBP1-51904) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-STAP2 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	55620
Gene Symbol	STAP2
Species	Human
Specificity/Sensitivity	This antibody is expected to recognize both reported isoforms (NP_060190.2; NP_001013863.1).
Immunogen	Peptide with sequence C-LQKKLEKRRALEH corresponding to C-Terminus according to NP_060190.2, NP_001013863.1.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Peptide ELISA
Recommended Dilutions	Western Blot 0.3 - 1 ug/mL, Immunohistochemistry-Paraffin 3.75 ug/mL, Peptide ELISA Detection limit 1:128000
Application Notes	WB: Approx. 49 kDa band observed in human breast lysates (calculated MW of 49.6 kDa band according to NP_060190.2).



Images


Western Blot: STAP2 Antibody [NBP1-51904] - Analysis of Human Breast lysate (35 ug protein in RIPA buffer). Antibody at 0.3 ug/mL. Detected by chemiluminescence.

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa

25kDa
20kDa

15kDa

Immunohistochemistry-Paraffin: STAP2 Antibody [NBP1-51904] - Staining of paraffin embedded Human Small Intestine. Antigen at 3.75 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Publications

Sekine Y, Ikeda O, Tsuji S et al. Signal-transducing adaptor protein-2 regulates stromal cell-derived factor-1 alpha-induced chemotaxis in T cells J Immunol 2009-12-15 [PMID: 19933863]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-51904

HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control
NBP1-88586PEP	STAP2 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-51904

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



