

Product Datasheet

TXNIP Antibody (JY2) - Azide and BSA Free NBP1-54578

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 3 Publications: 38

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-54578

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-54578



NBP1-54578

TXNIP Antibody (JY2) - Azide and BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JY2
Preservative	No Preservative
Isotype	IgG1
Purity	Protein A or G purified
Buffer	PBS


Product Description	
Description	Novus Biologicals Knockout (KO) Validated Mouse TXNIP Antibody (JY2) - Azide and BSA Free (NBP1-54578) is a monoclonal antibody validated for use in IHC, WB, Flow, Simple Western and IP. Anti-TXNIP Antibody: Cited in 33 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	10628
Gene Symbol	TXNIP
Species	Human, Mouse, Rat
Reactivity Notes	Use in Mouse reported in scientific literature (PMID:34576095). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Immunogen	Human recombinant TXNIP

Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation, Proximity Ligation Assay, CyTOF-ready, Knockout Validated
Recommended Dilutions	Western Blot 1 ug/ml, Simple Western 1:500, Flow Cytometry 2-5 ug/0.1x10 ⁶ cells, Immunohistochemistry, Immunoprecipitation 2ug/200ul of cell extract from 5x10 ⁶ cells, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen reported by customer review, Proximity Ligation Assay, CyTOF-ready, Knockout Validated reported in scientific literature (Brocker et al)




Images


Immunohistochemistry-Frozen: TXNIP Antibody (JY2) [NBP1-54578] - TXNIP in mouse brain. Antibody dilution: 1:100, incubated overnight in 5%BSA + 0.2% Triton X 100. Image from verified customer review.




Western Blot: TXNIP Antibody (JY2) [NBP1-54578] - TXNIP expression in Raji (1), K562 (2), KG1 (3), MRC5 (4), IC2Tr (5), HEL (6), P19 (7) and WR19L (8).




Western Blot: TXNIP Antibody (JY2) - Azide and BSA Free [NBP1-54578] - Analysis of TXNIP (NBP1-54578), ACTB, CASP1 (cleaved), and IL1B (cleaved) protein in livers from mice treated with WY-14643 for 48 h. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33203882/>) licensed under a CC-BY license.




Flow Cytometry: TXNIP Antibody (JY2) - Azide and BSA Free [NBP1-54578] - An intracellular stain was performed on THP-1 cells with TXNIP Antibody (JY2) NBP1-54578AF488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.




Western Blot: TXNIP Antibody (JY2) [NBP1-54578] - Hepa1 lysates probed with anti-TXNIP (JY2) at 1:2K. Cells were transfected with either empty vector or mouse TXNIP expression vector (MYC/DDK tagged). 30ug protein per lane. Image from verified customer review.




Western Blot: TXNIP Antibody (JY2) [NBP1-54578] - AMPK mediates anti-pyrototic effects of Exendin-4. (a) ROS determination by FACS. The left arrow indicated the ROS-negative population and the right pointed the positive. (b) Transcription activity of TXNIP in cardiomyocytes. (c) Western blot. (d) RNA silencing of TXNIP in cardiomyocytes. (e) IL-1 ELISA with TXNIP RNAi. (f) Caspase-1 activity assay with TXNIP RNAi. (g) Quantification of pAMPK/AMPK with CC treatment. (h) IL-1 ELISA with CC treatment. (i) Caspase-1 activity assay with CC treatment. Values are the mean SEM of 3 samples per group. 0.05, 0.01, 0.005, and 0.001. Exendin-4 Protects against Hyperglycemia-Induced Cardiomyocyte Pyroptosis via the AMPK-TXNIP Pathway. *J Diabetes Res* (2019)




Immunohistochemistry-Paraffin: TXNIP Antibody (JY2) [NBP1-54578] - Formalin fixed and paraffin embedded tissue section of mouse kidney using TXNIP antibody (clone JY2) at 1:250 dilution. The signal was developed using HRP-conjugated secondary antibody and DAB reagent which followed counterstaining of the cell nuclei with hematoxylin. This TXNIP antibody generated a diffused cytoplasmic signal in the cells of various tubules and glomeruli.




Immunohistochemistry-Paraffin: TXNIP Antibody (JY2) [NBP1-54578] - Formalin fixed and paraffin embedded tissue section of mouse kidney using TXNIP antibody (clone JY2) at 1:100 dilution. The signal was developed using HRP-conjugated secondary antibody and DAB reagent which followed counterstaining of the cell nuclei with hematoxylin. This TXNIP antibody generated a diffused cytoplasmic signal in the cells of various tubules with negligible staining in glomeruli.




Flow Cytometry: TXNIP Antibody (JY2) [NBP1-54578] - An intracellular stain was performed on RH-30 cells with TXNIP (JY2) Antibody NBP1-54578F (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to FITC.



Immunoprecipitation: TXNIP Antibody (JY2) [NBP1-54578] - Txnip/VDUP1 from Raji with NBP1-54578 (1) or mouse IgG1 (2).



Western Blot: TXNIP Antibody (JY2) - Azide and BSA Free [NBP1-54578] - AMPK mediates anti-pyrototic effects of Exendin-4. (a) ROS determination by FACS. The left arrow indicated the ROS-negative population & the right pointed the positive. (b) Transcription activity of TXNIP in cardiomyocytes. (c) Western blot. (d) RNA silencing of TXNIP in cardiomyocytes. (e) IL-1 β ELISA with TXNIP RNAi. (f) Caspase-1 activity assay with TXNIP RNAi. (g) Quantification of pAMPK/AMPK with CC treatment. (h) IL-1 β ELISA with CC treatment. (i) Caspase-1 activity assay with CC treatment. Values are the mean \pm SEM of 3 samples per group. \square $p < 0.05$, $\square\square$ $p < 0.01$, $\square\square\square$ $p < 0.005$, & $\square\square\square\square$ $p < 0.001$. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31886288>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Ismael S, Nasoohi S, Yoo A et al. Verapamil as an Adjunct Therapy to Reduce tPA Toxicity in Hyperglycemic Stroke: Implication of TXNIP/NLRP3 Inflammasome Molecular Neurobiology 2021-08-01 [PMID: 33847912]

Ismael S, Nasoohi S, Yoo A et al. Tissue Plasminogen Activator Promotes TXNIP-NLRP3 Inflammasome Activation after Hyperglycemic Stroke in Mice Molecular Neurobiology 2020-06-01 [PMID: 32172516]

Lim JO, Lee SJ, Kim WI et al. Melatonin Alleviates Silica Nanoparticle-Induced Lung Inflammation via Thioredoxin-Interacting Protein Downregulation Antioxidants (Basel) 2021-11-04 [PMID: 34829636]

Kim, J;Lim, J;Yoo, ID;Park, S;Moon, JS; TXNIP contributes to induction of pro-inflammatory phenotype and caspase-3 activation in astrocytes during Alzheimer's diseases Redox biology 2023-05-06 [PMID: 37172394] (Immunohistochemistry-Paraffin, Human)

Singh G Role of Glucose-induced Transcription Factor Signalling and Mitochondrial Epigenetics in Stress Tolerant Wood Frog, *Rana sylvatica* Thesis 2022-12-16 (Immunoprecipitation, Western Blot, Amphibian)

Details:
Wood Frog

Singh G, Storey KB TXNIP shuttling - a key molecular link in regulating inflammation and mitochondrial dysfunction in freeze tolerant wood frogs Gene 2023-01-07 [PMID: 36627089] (WB, Amphibian)

Patrick Devlin Inflammatory Response Following Hemorrhagic Stroke: The Role of Cytokines University of Tennessee Health Science Center 2022-11-07 (WB, Mouse)

Ismael S, Patrick D, Salman M et al. Verapamil inhibits TXNIP-NLRP3 inflammasome activation and preserves functional recovery after intracerebral hemorrhage in mice Neurochemistry international 2022-10-14 [PMID: 36244583] (WB, IF/IHC, Mouse)

Details:
Dilution used in WB 1:1000, in IHC 1:100

Girdhar K, Thakur S, Gaur P et al. Design, synthesis, and biological evaluation of a small molecule oral agonist of the glucagon-like-peptide-1 receptor Journal of Biological Chemistry 2022-04-01 [PMID: 35378127] (WB, Mouse)

Tsubaki H, Mendsaikhan A, Buyandelger U et al. Localization of Thioredoxin-Interacting Protein in Aging and Alzheimer's Disease Brains NeuroSci 2022-03-31 (IF/IHC, WB, Human)

Salman M, Ismael S, Lexiao L Et al. Acute Hyperglycemia Exacerbates Hemorrhagic Transformation after Embolic Stroke and Reperfusion with tPA: A Possible Role of TXNIP-NLRP3 Inflammasome Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association 2021-11-27 [PMID: 34847489] (WB, Mouse)

Salman M, Ismael S, Li L Et al. Endothelial Thioredoxin-Interacting Protein Depletion Reduces Hemorrhagic Transformation in Hyperglycemic Mice after Embolic Stroke and Thrombolytic Therapy Pharmaceuticals (Basel, Switzerland) 2021-09-27 [PMID: 34681207] (WB, Mouse)

More publications at <http://www.novusbio.com/NBP1-54578>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-54578

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP1-84784PEP	TXNIP Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-54578

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

