Product Datasheet

LAIR1 Antibody NBP1-68845

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-68845

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-68845



NBP1-68845

LAIR1 Antibody

Product Information 0.1 mg Concentration 0.5 mg/ml Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western Blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour 1HC:	LAIK I Antibody		
Concentration Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot :1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western Blot : Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Product Information		
Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Unit Size	0.1 mg	
Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Concentration	0.5 mg/ml	
Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western Blot Approx 28kDa band observed in Human Lymph Node Iysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Storage	Store at -20C. Avoid freeze-thaw cycles.	
Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western Blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Clonality	Polyclonal	
Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes (calculated MW of 29.8kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Preservative	0.02% Sodium Azide	
Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes (calculated MW of 29.8kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Isotype	IgG	
Product Description Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Purity	Immunogen affinity purified	
Novus Biologicals Goat LAIR1 Antibody (NBP1-68845) is a polyclonal antibody validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA	
validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 3903 Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Product Description		
Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Description	validated for use in IHC, WB and ELISA. All Novus Biologicals antibodies are	
Gene Symbol LAIR1 Species Human Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Host	Goat	
Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Gene ID	3903	
Specificity/Sensitivity This antibody is expected to recognize both isoforms (NP_002278.1; NP_068352.1). Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Gene Symbol	LAIR1	
Immunogen Peptide with sequence C-NDTEDVSQASPSE corresponding to internal region according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Species	Human	
according to NP_002278.1, NP_068352.1. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Specificity/Sensitivity		
Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Application Notes Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:1000 Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Immunogen		
Recommended Dilutions Western Blot 1:1000, Immunohistochemistry 1-3ug/ml, Immunohistochemistry- Paraffin, Peptide ELISA Detection limit 1:1000 Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Product Application Details		
Application Notes Western blot: Approx 28kDa band observed in Human Lymph Node lysates (calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Applications		
(calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	Recommended Dilutions		
	Application Notes	(calculated MW of 29.8kDa according to NP_068352.1). Primary incubation was 1 hour	



	1 ago 2 01 1 V.20.1 opaaloa 0/0/2020
Images	
Western Blot: LAIR1 Antibody [NBP1-68845] - Analysis of LAIR1 in Human Lymph Node lysate (35ug protein in RIPA buffer) using NBP1-68845 at 2ug/ml. Primary incubation was 1 hour. Detected by chemiluminescence.	250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa
Immunohistochemistry-Paraffin: LAIR1 Antibody [NBP1-68845] - 3.75ug/ml staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.	X



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP1-68845

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

NBP1-84590PEP LAIR1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-68845

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



