Product Datasheet

SLC22A3 Antibody - BSA Free NBP1-69368

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-69368

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-69368



NBP1-69368


SLC22A3 Antibody - BSA Free

| Product Information Unit Size 100 ul Concentration 0.5 mg/ml Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoarnine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Publications Western Blot, 1.0 ug/ml, Immunohistochemistry.Paraffin Recommended Dilutions Western Blot, 1.0 ug/ml, Immunohistochemistry 1:10-1:500, Immunohistochemistry.Paraffin 1:10-1:500 | - | |
|---|-----------------------------|---|
| Concentration 0.5 mg/ml Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Species expense Product Application Details Vestern Blot, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry, 1:10-1:500, | Product Information | |
| Storage Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (straneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Ppeduct Application Details Yeptide sequence for this immunogen was taken from within the described region. Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin | Unit Size | 100 ul |
| cycles.ClonalityPolyclonalPreservative0.09% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS, 2% SucroseTarget Molecular Weight61 kDaProduct DescriptionThe addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution.HostRabbitGene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Concentration | 0.5 mg/ml |
| Preservative 0.09% Sodium Azide Isotype IgG Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. Peptide sequence mas selected from the N terminal of SLC22A3. < | Storage | |
| Isotype IgG Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region. Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Clonality | Polyclonal |
| Purity Affinity purified Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence may be added region. Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Preservative | 0.09% Sodium Azide |
| Buffer PBS, 2% Sucrose Target Molecular Weight 61 kDa Product Description The addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution. Host Rabbit Gene ID 6581 Gene Symbol SLC22A3 Species Human Immunogen Synthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence pDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region. Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Isotype | IgG |
| Target Molecular Weight61 kDaProduct DescriptionInterventionDescriptionThe addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution.HostRabbitGene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Purity | Affinity purified |
| Product DescriptionDescriptionThe addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution.HostRabbitGene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Buffer | PBS, 2% Sucrose |
| DescriptionThe addition of 50% glycerol is optional for those storing this antibody at -20C and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution.HostRabbitGene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Target Molecular Weight | 61 kDa |
| and not aliquoting smaller units. However, please note that glycerol may interrupt some downstream antibody applications and should be added with caution.HostRabbitGene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application Details ApplicationsWestern Blot, Immunohistochemistry, Immunohistochemistry-Paraffin Western Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Product Description | |
| Gene ID6581Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Description | and not aliquoting smaller units. However, please note that glycerol may interrupt |
| Gene SymbolSLC22A3SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application DetailsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Host | Rabbit |
| SpeciesHumanImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application Details ApplicationsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Gene ID | 6581 |
| ImmunogenSynthetic peptides corresponding to SLC22A3(solute carrier family 22 (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the describedProduct Application Details ApplicationsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Gene Symbol | SLC22A3 |
| (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described region.Product Application Details ApplicationsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Species | Human |
| ApplicationsWestern Blot, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Immunogen | (extraneuronal monoamine transporter), member 3) The peptide sequence was selected from the N terminal of SLC22A3. Peptide sequence PDHYWCRGPSAAALAERCGWSPEEEWNRTAPASRGPEPPERRGRCQRYLL The peptide sequence for this immunogen was taken from within the described |
| Recommended Dilutions Western Blot 1.0 ug/ml, Immunohistochemistry 1:10-1:500, | Product Application Details | |
| | Applications | Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin |
| | Recommended Dilutions | |




Images

Western Blot: SLC22A3 Antibody [NBP1-69368] - This Anti-SLC22A3 antibody was used in Western Blot of HepG2 tissue lysate at a concentration of 0.5ug/ml.



Immunohistochemistry: SLC22A3 Antibody [NBP1-69368] - Human kidney







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-69368

| NBP1-83976PEP | SLC22A3 Recombinant Protein Antigen |
|---------------|---|
| NBP2-24891 | Rabbit IgG Isotype Control |
| NB7160 | Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP] |
| HAF008 | Goat anti-Rabbit IgG Secondary Antibody [HRP] |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-69368

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

