# **Product Datasheet**

# Goat anti-Hamster IgG (H+L) Secondary Antibody [Allophycocyanin/Cy7] (Pre-adsorbed) NBP1-71730APCCY7

Unit Size: 500 ul

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-71730APCCY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-71730APCCY7



# **NBP1-71730APCCY7**

Goat anti-Hamster IgG (H+L) Secondary Antibody [Allophycocyanin/Cy7] (Pre-adsorbed)

| Goat anti-Hamster IgG (H+L) Secondary Antibody [Allophycocyanin/Cy7] (Pre-adsorbed) |  |
|---|--|
| Product Information   |  |
| Unit Size   | 500 ul   |
| Concentration   | Please see the vial label for concentration. If unlisted please contact technical services.  |
| Storage   | Store at 4C in the dark. Do not freeze.  |
| Clonality   | Polyclonal   |
| Preservative  | 0.05% Sodium Azide   |
| Isotype   | IgG  |
| Conjugate   | Allophycocyanin/Cy7  |
| Purity  | Multi-step   |
| Buffer  | PBS  |
| Product Description   |  |
| Host  | Goat   |
| Species   | Hamster  |
| Reactivity Notes  | Cricetulus migratorius (Armenian Hamster); Melanochromis auratus (Golden Syrian Hamster)   |
| Specificity/Sensitivity   | This antibody was pre-adsorbed against Mouse and Rat Serum Proteins.   |
|   | No reaction was observed against Mouse or Rat serum proteins. Specificity was confirmed using ELISA at less than 1% cross reactivity.  |
| Immunogen   | Armenian and Golden Syrian Hamster IgG, whole molecule   |
| Product Application Details   |  |
| Applications  | Western Blot, Dot Blot, ELISA, Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry   |
| Recommended Dilutions   | Western Blot, ELISA, Immunohistochemistry, Immunocytochemistry/<br>Immunofluorescence, Dot Blot, Fluorophore-linked immunosorbent assay  |
| Application Notes   | Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions. |





## **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

# **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-71730APCCY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

