

# Product Datasheet

## Vanilloid R1/TRPV1 Antibody - BSA Free NBP1-71774

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1 Publications: 12**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-71774](http://www.novusbio.com/NBP1-71774)

Updated 12/20/2023 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-71774](http://www.novusbio.com/reviews/destination/NBP1-71774)



**NBP1-71774**

Vanilloid R1/TRPV1 Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description	
Host	Rabbit
Gene ID	7442
Gene Symbol	TRPV1
Species	Human, Mouse
Immunogen	A genomic peptide made to a C-terminal region of the human TRPV1 protein (within residues 725-839). [Swiss-Prot# Q8NER1]
Notes	Manufactured by Genomic Antibody Technology™. GAT <a href="#">FAQs</a>


Product Application Details	
Applications	Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry 1:200, Immunocytochemistry/ Immunofluorescence 1:50-1:100, Immunohistochemistry-Paraffin 1:200
Application Notes	This TRPV1 antibody is useful for ICC/IF and IHC-paraffin embedded tissues. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.

**Images**


Immunohistochemistry: Vanilloid R1/TRPV1 Antibody [NBP1-71774] - The Vanilloid R1/TRPV1+ and Vanilloid R1/TRPV1+CD4+ immune cells in patients with allergic rhinitis. Immunohistochemical staining of Vanilloid R1/TRPV1 (single staining) and CD4/Vanilloid R1/TRPV1 (double staining) in patients with allergic rhinitis compared with control subjects. Image collected and cropped by CiteAb from the following publication (<https://www.oncotarget.com/fulltext/6653>), licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: Vanilloid R1/TRPV1 Antibody [NBP1-71774] - TRPV1 antibody was tested at 1:50 in HeLa cells with FITC (green). Nuclei were counterstained with DAPI (blue).




Immunohistochemistry: Vanilloid R1/TRPV1 Antibody [NBP1-71774] - Analysis of TRPV1 in mouse dorsal root ganglion using DAB with hematoxylin counterstain.



Immunohistochemistry-Paraffin: Vanilloid R1/TRPV1 Antibody [NBP1-71774] - Human stomach tissue. Positive staining can be observed in gastric neurons. Image from verified customer review.



The TRPV1+ and TRPV1+CD4+ immune cells in patients with allergic rhinitis. Immunohistochemical staining of TRPV1 (single staining) and CD4/TRPV1 (double staining) in patients with allergic rhinitis compared with control subjects B. Counts of TRPV1 + cells and TRPV1+ CD4+ cells in each groups. The statistical P values are presented as \* (P < 0.05), and \*\* (P < 0.01).



## Publications

Meilinn Tram, Tarek Ibrahim, Anahit Hovhannisyanyan, Armen Akopian, Shivani Ruparel Lingual innervation in male and female marmosets *Neurobiology of Pain* 2023-05-31 [PMID: 38099285]

Warfield R, Robinson JA, Podgorski RM et al. Neuroinflammation in the Dorsal Root Ganglia and Dorsal Horn Contributes to Persistence of Nociceptor Sensitization in SIV-Infected Antiretroviral Therapy-Treated Macaques *The American journal of pathology* 2023-09-19 [PMID: 37734588] (IHC)

Tram M, Ibrahim T, Hovhannisyanyan A et al. Lingual innervation in male and female marmosets *Neurobiology of Pain* 2023-08-01 (IHC-Fr, Primate)

### Details:

Species-marmoset (*Callithrix jacchus*)

Kumar V, Kingsley D, Madhurakkat Perikamana S et al. Self-assembled innervated vasculature-on-a-chip to study nociception *Biofabrication* 2023-04-13 [PMID: 36996841] (WB)

Hovhannisyanyan AH, Lindquist KA, Belugin S et al. Sensory innervation of masseter, temporal and lateral pterygoid muscles in common marmosets *bioRxiv : the preprint server for biology* 2023-02-12 [PMID: 36798270] (IHC-Fr, Marmoset, Primate)

### Details:

Dilution used in IHC-Fr 1:200

Ilhan HD, Unal B, Ayaz Y, Erin N Changes in TRPV1 Expression as Well as Substance P and Vasoactive Intestinal Peptide Levels Are Associated with Recurrence of Pterygium *International journal of molecular sciences* 2022-12-10 [PMID: 36555331] (IHC-P, Human)

### Details:

Dilution used in IHC-P 1:100

Moutafidi A, Gatzounis G, Zolota V, Assimakopoulou M Heat shock factor 1 in brain tumors: a link with transient receptor potential channels TRPV1 and TRPA1 *Journal of molecular histology* 2021-09-30 [PMID: 34591198]

Bi X, Xu Y, Li T et al. Chronic stress augments esophageal inflammation, and alters the expression of transient receptor potential vanilloid 1 and protease-activated receptor 2 in a murine model *Mol Med Rep* 2019-06-01 [PMID: 31059059]

Rizopoulos T, Papadaki-Petrou H, Assimakopoulou M. Expression Profiling of the Transient Receptor Potential Vanilloid (TRPV) Channels 1, 2, 3 and 4 in Mucosal Epithelium of Human Ulcerative Colitis. *Cells*. 2018-06-15 [PMID: 29914124] (IHC-P, Human)

Assimakopoulou M, Pagoulatos D, Nterma P, Pharmakakis N. Immunolocalization of cannabinoid receptor type 1 and CB2 cannabinoid receptors, and transient receptor potential vanilloid channels in pterygium *Mol Med Rep* 2017-08-14 [PMID: 28849159] (Human)

Hidaka T, Ogawa E, Kobayashi EH et al. The aryl hydrocarbon receptor AhR links atopic dermatitis and air pollution via induction of the neurotrophic factor artemin. *Nat. Immunol.* 2016-11-21 [PMID: 27869817] (Mouse)

### Details:

This citation used the DyLight 488 version of this antibody.

Samivel R, Kim DW, Son HR et al. The role of TRPV1 in the CD4+ T cell-mediated inflammatory response of allergic rhinitis. *Oncotarget*. 2016-01-05 [PMID: 26700618] (IF/IHC, Human, Mouse)

## Procedures

### Immunohistochemistry-Paraffin protocol for TRPV1 Antibody (NBP1-71774)

Vanilloid R1/TRPV1 Antibody:

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

### Immunocytochemistry/Immunofluorescence Protocol for TRPV1 Antibody (NBP1-71774)

Vanilloid R1/TRPV1 Antibody:

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures."



### Novus Biologicals USA

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### Bio-Techne Canada

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### Bio-Techne Ltd

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### General Contact Information

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### Products Related to NBP1-71774

---

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-71774G	Vanilloid R1/TRPV1 Antibody [DyLight 488]

---

### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-71774](http://www.novusbio.com/reviews/submit/NBP1-71774)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

