Product Datasheet

Goat anti-Human IgA Heavy Chain Secondary Antibody [HRP] (Pre-adsorbed) NBP1-74913

Unit Size: 1 mg

Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.

www.novusbio.com

technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-74913

Updated 10/30/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-74913



NBP1-74913

Goat anti-Human IgA Heavy Chain Secondary Antibody [HRP] (Pre-adsorbed)

Unit Size1 mgConcentrationLYOPH mg/mlStorageStore lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.ClonalityPolyclonalPreservative0.1% KathonReconstitution InstructionsRehydrate with 1.1 ml of deionized water and let stand 30 minutes at room recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution recovery.) Centrifuge to remove any particulates.IsotypeIgGConjugateHRPPurityAffinity purifiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionStore lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins.Specificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended DilutionsNotionWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry 1:500, ILISO0, Immunohistochemistry-Paraffin<	Product Information	
StorageStore lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.ClonalityPolyclonalPreservative0.1% KathonReconstitution InstructionsRehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recover.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.IsotypeIgGConjugateHRPPurityAffinity purfiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protaese/IgG freeProduct DescriptionPurity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily.Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins.Specificity/SensitivityBased on IEP, nis autibody reacts with heavy alpha chains on human IgA. This artibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-	Unit Size	1 mg
glycerol, aliquot contents and freeze at -20C or below for long term storage.ClonalityPolyclonalPreservative0.1% KathonReconstitution InstructionsRehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.IsotypeIgGConjugateHRPPurityAffinity purfiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionDescriptionStore lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin serum proteins from bovine, mouse, or rabbitSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulins serum proteins from bovine, mouse, or rabbitSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulins serum proteins from bovine, mouse, or rabbitProduct Application DetailsPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, 1:200-1:5000, Immunohistochemistry-ParaffinReactivity NotesWestern Blot, 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-	Concentration	LYOPH mg/ml
Preservative 0.1% Kathon Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate HRP Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Purity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animats of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins. from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins. Immunogen Purified human IgA, (Alpha chain) Product Application Details Western Blot, 1:200-1:5000, Immunohistochemistry-Paraffin	Storage	
Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate HRP Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Purity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulin serum proteins. Specificity/Sensitivity Based on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins. Immunogen Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500	Clonality	Polyclonal
temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.IsotypeIgGConjugateHRPPurityAffinity purifiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionPurity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily.BostGoatStore lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry. Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot, 1:200-1:5000, ILISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Preservative	0.1% Kathon
ConjugateHRPPurityAffinity purifiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionDescriptionPurity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry. Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Reconstitution Instructions	temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution
PurityAffinity purifiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionPurity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Isotype	lgG
Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Purity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins. Immunogen Purified human IgA, (Alpha chain) Product Application Details Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1:200-1:5000, Immunohistochemistry 1:500-1:5000,	Conjugate	HRP
Protease/IgG free Product Description Description Purity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins. Immunogen Purified human IgA, (Alpha chain) Product Application Details Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry -Paraffin	Purity	Affinity purified
DescriptionPurity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended DilutionsRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Buffer	
animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution, dilute with 50% glycerol and store at -20C as a liquid.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended DilutionsRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Product Description	
HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application DetailsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Description	animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily. Store lyophilized material at 2-8C. For long term storage after reconstitution,
Reactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum 	Host	Goat
immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy alpha chains on human IgA. This antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application Details ApplicationsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Species	Human
antibody has been pre-adsorbed against bovine, mouse, and rabbit serum proteins.ImmunogenPurified human IgA, (Alpha chain)Product Application Details ApplicationsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500- 1:5000, Immunohistochemistry-Paraffin	Reactivity Notes	immunoglobulins and light chains on all human immunoglobulins serum proteins
Product Application Details Applications Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry-Paraffin	Specificity/Sensitivity	antibody has been pre-adsorbed against bovine, mouse, and rabbit serum
ApplicationsWestern Blot, ELISA, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry-Paraffin	Immunogen	Purified human IgA, (Alpha chain)
Recommended Dilutions Western Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry-Paraffin	Product Application Details	
1:5000, Immunohistochemistry-Paraffin	Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin
Application NotesThis conjugate is suitable for all immunoassay applications.	Recommended Dilutions	
	Application Notes	This conjugate is suitable for all immunoassay applications.

Publications

Abu-Humaidan AHA, Ahmad FM, Awajan D et al. Anti-SARS-Cov-2 S-RBD IgG Formed after BNT162b2 Vaccination Can Bind C1q and Activate Complement Journal of immunology research 2022-12-17 [PMID: 36573216] (ELISA)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-74913

NBP3-20648-100ug

IgA Heavy Chain Antibody (IGHA/7567R)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-74913

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

