

Product Datasheet

Donkey anti-Rabbit IgG (H+L) Secondary Antibody [PE/Cy5.5] NBP1-75286PECY55

Unit Size: 500 ul

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-75286PECY55

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-75286PECY55



NBP1-75286PECY55

Donkey anti-Rabbit IgG (H+L) Secondary Antibody [PE/Cy5.5]

Product Information	
Unit Size	500 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark. Do not freeze.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Conjugate	PE/Cy5.5
Purity	Affinity purified
Buffer	PBS
Product Description	
Host	Donkey
Species	Rabbit
Reactivity Notes	Based on IEP, this antibody reacts with heavy (gamma) chains on rabbit IgG and light chains on all rabbit immunoglobulins.
Specificity/Sensitivity	Based on IEP, this Donkey anti-Rabbit IgG (H+L) Secondary Antibody (Pre-adsorbed) heavy gamma chains on rabbit IgG and light chains on all rabbit immunoglobulins. This antibody has been pre-adsorbed against bovine, chicken, goat, guinea pig, hamster, horse, human, mouse, rat or sheep
Immunogen	This Donkey anti-Rabbit IgG (H+L) Secondary Antibody (Pre-adsorbed) was developed against purified rabbit IgG (H&L).
Product Application Details	
Applications	Western Blot, ELISA, Flow Cytometry, Immunoassay, Immunocytochemistry/ Immunofluorescence, Immunodiffusion, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot, Flow Cytometry, ELISA, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Immunohistochemistry-Frozen, Immunodiffusion, Immunoassay
Application Notes	Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any questions.

Publications

Florian Hamberger, Young-Seon Mederacke, Ingmar Mederacke An inducible model for genetic manipulation and fate-tracing of PDGFR β -expressing fibrogenic cells in the liver Scientific Reports 2023-05-05 [PMID: 37147343]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-75286PECY55

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

