Product Datasheet

Goat anti-Human IgA Heavy Chain Secondary Antibody [DyLight 550] (Pre-adsorbed) NBP1-75759

Unit Size: 1 mg

Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-75759

Updated 10/30/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-75759



NBP1-75759

Goat anti-Human IgA Heavy Chain Secondary Antibody [DyLight 550] (Pre-adsorbed)

Unit Size 1 mg Concentration LVOPH mg/ml Storage Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage. Clonality Polyclonal Preservative 0.05% Sodium Azide Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfiled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Strange after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins. from bovine, mouse, or rabbit Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins. from bovine, mouse, or rabbit Specificity/Sensi	Product Information	
Storage Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage. Clonality Polyclonal Preservative 0.05% Sodium Azide Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product bescription Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human	Unit Size	1 mg
glycerol, aliquot contents and freeze at -20C or below for long term storage. Clonality Polyclonal Preservative 0.05% Sodium Azide Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Joretase/IgG free Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered verterinarian. DyLight 550 (Ex = 550 nm). After reconstituted, prepare fresh working dilution daily Product bescription Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered verterinarian. DyLight 550 (Ex = 550 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is obs	Concentration	LYOPH mg/ml
Preservative 0.05% Sodium Azide Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins form bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries	Storage	
Reconstitution Instructions Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily. Isotype IgG Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Description Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product bescription Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibo	Clonality	Polyclonal
temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.IsotypeIgGConjugateDyLight 550PurityAffinity purifiedBuffer10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG freeProduct DescriptionAffinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Preservative	0.05% Sodium Azide
Conjugate DyLight 550 Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunomicroscopy Recommended Dilutions Flow Cytometry 1:20 - 1:2000	Reconstitution Instructions	temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution
Purity Affinity purified Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunomicroscopy Applications Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Isotype	IgG
Buffer 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free Product Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunomicroscopy 1:20 - 1:2000	Conjugate	DyLight 550
Protease/IgG free Product Description Description Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C. Host Goat Species Human Reactivity Notes Based on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbit Specificity/Sensitivity Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Flow Cytometry, Immunomicroscopy Applications Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Purity	Affinity purified
DescriptionAffinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution dailyProduct is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Buffer	
obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution dailyProduct is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C.HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Product Description	
HostGoatSpeciesHumanReactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Description	obtained from healthy animals of US origin and under the care of a registered veterinarian. DyLight 550 (Ex = 550 nm; Em = 576 nm). After reconstituted, prepare fresh working dilution daily Product is stable for up to 4 weeks at 2-8C after rehydration. For extended
Reactivity NotesBased on IEP, no reactivity is observed to non-immunoglobulin human serum immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Host	
immunoglobulins and light chains on all human immunoglobulins serum proteins from bovine, mouse, or rabbitSpecificity/SensitivityBased on IEP, this antibody reacts with heavy (alpha) chains on human IgAImmunogenPurified human IgA, (Alpha chain)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Species	Human
Immunogen Purified human IgA, (Alpha chain) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Flow Cytometry, Immunomicroscopy Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Reactivity Notes	immunoglobulins and light chains on all human immunoglobulins serum proteins
NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsFlow Cytometry, ImmunomicroscopyApplicationsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Specificity/Sensitivity	Based on IEP, this antibody reacts with heavy (alpha) chains on human IgA
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsApplicationsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Immunogen	Purified human IgA, (Alpha chain)
Product Application Details Applications Flow Cytometry, Immunomicroscopy Recommended Dilutions Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Notes	
ApplicationsFlow Cytometry, ImmunomicroscopyRecommended DilutionsFlow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000		DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Recommended Dilutions Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000	Product Application Details	
	Applications	Flow Cytometry, Immunomicroscopy
Application NotesThis conjugate is suitable for immunomicroscopy and flow cytometry.	Recommended Dilutions	Flow Cytometry 1:20 - 1:2000, Immunomicroscopy 1:20 - 1:2000
	Application Notes	This conjugate is suitable for immunomicroscopy and flow cytometry.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-75759

NBP3-20648-100ug

IgA Heavy Chain Antibody (IGHA/7567R)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-75759

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

