

Product Datasheet

TRAILR4/TNFRSF10D/DcR2 Antibody - BSA Free NBP1-76985

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 20


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-76985

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-76985



NBP1-76985


TRAILR4/TNFRSF10D/DcR2 Antibody - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS
Target Molecular Weight	36 kDa
Product Description	
Description	Novus Biologicals Rabbit TRAILR4/TNFRSF10D/DcR2 Antibody - BSA Free (NBP1-76985) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow and ICC/IF. Anti-TRAILR4/TNFRSF10D/DcR2 Antibody: Cited in 20 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	8793
Gene Symbol	TNFRSF10D
Species	Human, Mouse, Rat
Immunogen	Antibody was raised against a 15 amino acid peptide near the center of human DcR2. The immunogen is located within amino acids 230 - 280 of DcR2. Amino Acid Sequence: GGPERVHRVLFRRRS
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Flow (Cell Surface), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1 ug/ml, Flow Cytometry, ELISA 1:100-1:2000, Immunohistochemistry 2.5 ug/ml, Immunocytochemistry/ Immunofluorescence 10 -20 ug/ml, Immunohistochemistry-Paraffin 2.5 ug/ml, Flow (Cell Surface)
Application Notes	Use in FLOW reported in scientific literature (PMID: 12795746). Use in Flow-cell surface reported in scientific literature (PMID: 12173037).




Images


Western Blot: TRAILR4/TNFRSF10D/DcR2 Antibody [NBP1-76985] - Analysis of 50 ug of whole cell lysate from HeLa cells with anti-DcR2 at 1:1000 dilution.



Immunocytochemistry/Immunofluorescence: TRAILR4/TNFRSF10D/DcR2 Antibody [NBP1-76985] - Staining of HeLa cells using DcR2 antibody at 10ug/ml.




Immunohistochemistry: TRAILR4/TNFRSF10D/DcR2 Antibody - BSA Free [NBP1-76985] - Immunohistochemistry of TRAILR4/TNFRSF10D/DcR2 in mouse kidney node tissue with TRAILR4/TNFRSF10D/DcR2 antibody at 2.5 ug/ml.



Immunocytochemistry/ Immunofluorescence: TRAILR4/TNFRSF10D/DcR2 Antibody - BSA Free [NBP1-76985] - Immunofluorescence of TRAILR4/TNFRSF10D/DcR2 in mouse kidney tissue with TRAILR4/TNFRSF10D/DcR2 antibody at 20 ug/ml.

Green: TRAILR4/TNFRSF10D/DcR2 Antibody
Blue: DAPI staining



Publications

Fang Y, Lu J, Wang X et al. HIF-1 alpha Mediates TRAIL-Induced Neuronal Apoptosis via Regulating DcR1 Expression Following Traumatic Brain Injury Front Cell Neurosci 2020-08-03 [PMID: 32848609] (WB, Rat)

Koyama S. Flow cytometric measurement of tumor necrosis factor-related apoptosis-inducing ligand and its receptors in gastric epithelium and infiltrating mucosal lymphocytes in *Helicobacter pylori*-associated gastritis. *J Gastroenterol Hepatol.* 2003-07-01 [PMID: 12795746] (FLOW)

Basile JR, Zacny V, Munger K. The cytokines tumor necrosis factor-alpha (TNF-alpha) and TNF-related apoptosis-inducing ligand differentially modulate proliferation and apoptotic pathways in human keratinocytes expressing the human papillomavirus-16 E7 oncoprotein. *J Biol Chem.* 2001-06-22 [PMID: 11306566] (WB, Human)

Details:

WB DR5/Apo2/TRAIL-R2/TRAILR2/TRICK2/KILLER (IMG-120A) and DcR2/TRAIL-R4/TRUNDD (IMG-121). Human keratinocytes (HFK) cells expressing HPV-16 EF (human papillomavirus), Fig 6.

Mitsiades CS, Treon SP, Mitsiades N et al. TRAIL/Apo2L ligand selectively induces apoptosis and overcomes drug resistance in multiple myeloma: therapeutic applications. *Blood.* 2001-08-01 [PMID: 11468181]

Gupta SC, Francis SK, Nair MS et al. Azadirone, a Limonoid Tetranortriterpene, Induces Death Receptors and Sensitizes Human Cancer Cells to Tumor Necrosis Factor-related Apoptosis-inducing Ligand (TRAIL) through a p53 Protein-independent Mechanism: EVIDENCE FOR THE ROLE OF THE ROS-ERK-CHOP-DEATH RECEPTOR PATHWAY. *J Biol Chem* 2013-11-08 [PMID: 24078627] (WB, Human)

Mitsiades N, Poulaki V, Tseleni-Balafouta S et al. Thyroid carcinoma cells are resistant to FAS-mediated apoptosis but sensitive to tumor necrosis factor-related apoptosis-inducing ligand. *Cancer Res.* 2000-08-01 [PMID: 10945619]

Allen JE, Ferrini R, Dicker DT et al. Targeting TRAIL death receptor 4 with trivalent DR4 Atrimer complexes. *Mol Cancer Ther.* 2012-10-01 [PMID: 22802267]

Details:

Antibodies cited: 1. DR5 (IMG-120A): Flow (cell surface) 2. DR4 (IMG-141): Flow (cell surface, HCT116 cells), Supplementary Fig. 5. 3. DcR2 (IMG-121-1): WB (SW620 and HCT15 cells), Supplementary Fig. 4.

Lunghi P, Giuliani N, Mazzera L et al. Targeting MEK/MAPK signal transduction module potentiates ATO-induced apoptosis in multiple myeloma cells through multiple signaling pathways. *Blood.* 2008-09-15 [PMID: 18583568] (WB, Human)

Details:

1. p73 (IMG-246): WB (human myeloma cell lines (HMCLs) XG-6 and XG-1), Fig. 4Ci. 2. p73 (deltaNp73) [IMG-313A]: WB (HMCLs), Fig. 4Ai. 3. DcR2 (IMG-121): WB (HMCLs), Fig. 5A.

Fan QL, Zou WY, Song LH et al. Synergistic antitumor activity of TRAIL combined with chemotherapeutic agents in A549 cell lines in vitro and in vivo. *Cancer Chemother Pharmacol.* 2005-02-01 [PMID: 15290100]

Zhang HY, Wang HQ, Liu HM et al. Regulation of tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis by DJ-1 in thyroid cancer cells. *Endocr Relat Cancer.* 2008-06-01 [PMID: 18430896] (WB, Human)

Details:

WB (human thyroid cells).

Poulaki V, Mitsiades CS, Kotoula V et al. Regulation of Apo2L/tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis in thyroid carcinoma cells. *Am J Pathol.* 2002-08-01 [PMID: 12163389]

Nagane M, Shimizu S, Mori E et al. Predominant antitumor effects by fully human anti-TRAIL-receptor 2 (DR5) monoclonal antibodies in human glioma cells in vitro and in vivo. *Neuro Oncol.* 2010-07-01 [PMID: 20511188] (WB, Human)

Details:

Products cited: 1. DcR2 (IMG-121-1): WB (human glioma cell lines), Fig 4B. 2. DcR1 (IMG-245-1): WB (human glioma cell lines), Fig 4B.

More publications at <http://www.novusbio.com/NBP1-76985>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-76985

NBP1-76985PEP	TRAILR4/TNFRSF10D/DcR2 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-76985

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

