# **Product Datasheet**

# Lamin A + C R482W Antibody (5H8-B4) NBP1-77402

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-77402

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-77402



## NBP1-77402

Lamin A + C R482W Antibody (5H8-B4)


<b>Product Information</b>	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	5H8-B4
Preservative	0.1% Sodium Azide
Isotype	IgG1 Kappa
Purity	Unpurified
Buffer	Ascites
	·

Product Description	
Host	Mouse
Gene ID	4000
Gene Symbol	LMNA
Species	Human
Specificity/Sensitivity	Disease-associated point-mutant R482W of Lamin A + C
Immunogen	Hepatitis B Virus Capsid Antigen (HBcAg) Lamin A + C Ig-fold R482W fusion protein.

	protein.
<b>Product Application Details</b>	
Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000, ELISA 1:100-1:2000, Immunocytochemistry/ Immunofluorescence 1:10-1:2000, Immunoprecipitation 1:10-1:500
Application Notes	This Lamin A + C R482W (5H8-B4) point-mutant specific antibody is useful for Western blot, ELISA, Immunoprecipitation and Immunocytochemistry/Immunofluorescence. In WB bands can be seen at 74kDa (Lamin A) and 65 kDa (Lamin C).

## **Images**

Western Blot: Lamin A + C R482W Antibody (5H8-B4) [NBP1-77402] - Analysis of Lamin A + C R482W in HeLa cells transfected with Flagtagged Lamin A 1) wild type, 2) R482W and 3) R453W.





## **Publications**

Roblek M, Schuchner S, Huber V, Ollram K, Vlcek-Vesely S, Foisner R, Wehnert M, Ogris E. Monoclonal antibodies specific for disease-associated point-mutants: lamin A/C R453W and R482W. PLoS One. 5(5):e10604. 2010-05-13 [PMID: 20498701] (ICC/IF, IP, WB, Human)



#### **Procedures**

### Western Blot Protocol Specific for NBP1-77402: Lamin A + C R482W Antibody (5H8-B4)

Lamin A + C R482W Antibody (5H8-B4):

Western Blot Protocol

- 1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
- 2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
- 3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
- 4. Rinse the blot.
- 5. Block the membrane using standard blocking buffer for at least 1 hour.
- 6. Wash the membrane in wash buffer three times for 10 minutes each.
- 7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
- 8. Wash the membrane in wash buffer three times for 10 minutes each.
- 9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
- 10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
- 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- \*\*Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.
- \*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





## **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP1-77402**

**HAF007** Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

210-TA-005 TNF-alpha [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-77402

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

