Product Datasheet

ERp57/PDIA3 Antibody - BSA Free NBP1-84796

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-84796

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-84796




NBP1-84796

ERp57/PDIA3 Antibody - BSA Free

Product Information		
Unit Size	0.1 ml	
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.	
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Clonality	Polyclonal	
Preservative	0.02% Sodium Azide	
Isotype	IgG	
Purity	Immunogen affinity purified	
Buffer	PBS (pH 7.2) and 40% Glycerol	
Product Description		
Host	Rabbit	
Gene ID	2923	
Gene Symbol	PDIA3	
Species	Human, Mouse, Rat	
Marker	Endoplasmic Reticulum Marker	
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: PTLKIFRDGEEAGAYDGPRTADGIVSHLKKQAGPASVPLRTEEEFKKFISDKDA SIVGFFDDSFSEAHSEFLKAASNLRDNYRFAHTNVESLVNEYDDNGEGIILFRPS HLTNKFEDK	
Product Application Details		
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockdown Validated	
Recommended Dilutions	Western Blot 0.04 - 0.4 ug/ml, Immunohistochemistry 1:2500 - 1:5000, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:2500 - 1:5000, Knockdown Validated	
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.	

Images

Immunohistochemistry-Paraffin: ERp57/PDIA3 Antibody [NBP1-84796] -Staining in human thyroid gland and skeletal muscle tissues using NBP1 -84796 antibody. Corresponding PDIA3 RNA-seq data are presented for the same tissues.




PDIA3 in Thyroid gland

d PDIA3 in Skeletal muscle

```
RNA [TPM]
```

RNA [TPM] 20 40 60 80 100






www.novusbio.com



	Page 3 of 5 v.20.1 Updated 2/21/2025
Western Blot: ERp57/PDIA3 Antibody [NBP1-84796] - Analysis in mouse cell line NIH-3T3 and rat cell line NBT-II.	[kDa] 100 100 100 70 55 35 15 10 10
Immunohistochemistry-Paraffin: ERp57/PDIA3 Antibody [NBP1-84796] - Staining of human prostate shows moderate cytoplasmic positivity in glandular cells.	
Immunohistochemistry-Paraffin: ERp57/PDIA3 Antibody [NBP1-84796] - Staining of human skeletal muscle shows no cytoplasmic positivity in myocytes as expected.	
Immunohistochemistry-Paraffin: ERp57/PDIA3 Antibody [NBP1-84796] - Staining of human small intestine shows moderate to strong cytoplasmic positivity in glandular cells.	



Immunohistochemistry-Paraffin: ERp57/PDIA3 Antibody [NBP1-84796] - Staining of human thyroid gland shows strong cytoplasmic positivity in glandular cells.



Publications

Bechor E, Dahan I, Fradin T et al. The dehydrogenase region of the NADPH oxidase component Nox2 acts as a protein disulfide isomerase (PDI) resembling PDIA3 with a role in the binding of the activator protein p67phox. Front Chem 2015-01-01 [PMID: 25699251] (WB, Human)

Feng H, Li X, Chan V, Chen WN. Proteomics Based Identification of Cell Migration Related Proteins in HBV Expressing HepG2 Cells. PLoS One 2014-01-01 [PMID: 24763314] (WB)

Logan CV, Szabadkai G, Sharpe JA et al. Loss-of-function mutations in MICU1 cause a brain and muscle disorder linked to primary alterations in mitochondrial calcium signaling. Nat Genet 2014-02-01 [PMID: 24336167]

Racape M, Duong Van Huyen JP, Danger R et al. The involvement of SMILE/TMTC3 in endoplasmic reticulum stress response. PLoS One 2011-01-01 [PMID: 21603654]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-84796

NBP1-84796PEP	ERp57/PDIA3 Recombinant Protein Antigen
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-84796

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

