Product Datasheet NFkB2/NFkB p100 Antibody - BSA Free

NBP1-87760

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-87760

Updated 4/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-87760




NBP1-87760

NFkB2/NFkB p100 Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol
Product Description	
Host	Rabbit
Gene ID	4791
Gene Symbol	NFKB2
Species	Human
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: LLNAAQNTMEPPLTPPSPAGPGLSLGDTALQNLEQLLDGPEAQGSWAELAERL GLRSLVDTYRQTTSPSGSLLRSYELAGGDLAGLLEALSDMGLEEGVRLLRGPE TRDKLPSTEVKEDSAYGSQSVEQEA
Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockdown Validated
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Simple Western 1:30, Immunohistochemistry 1:200 - 1:500, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:500 - 1:1000, Knockdown Validated
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See <u>Simple Western Antibody Database</u> for Simple Western validation: Tested in RT4, separated by Size, antibody dilution of 1:30, apparent MW was 134 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.








Immunocytochemistry/Immunofluorescence: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human cell line A-431 shows localization to cytosol. Antibody staining is shown in green.






Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human appendix shows high expression.

Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human

testis.

Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human skeletal muscle shows low expression as expected.





Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human kidney.


Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human lymph node.


Immunohistochemistry-Paraffin: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Staining of human cerebral cortex.

Simple Western: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Simple Western lane view shows a specific band for NFKB2 in 0.2 mg/ml of RT4 lysate. This experiment was performed under reducing conditions using the 66-440 kDa separation system.


www.novusbio.com







Simple Western: NFkB2/NFkB p100 Antibody [NBP1-87760] - NFkB p100/p52 Antibody [NBP1-87760] - Electropherogram image(s) of corresponding Simple Western lane view. NFkB p100/p52 antibody was used at 1:30 dilution on RT4 lysate(s).



Publications

Xu W, Mo J, Ocak U et al. Activation of Melanocortin 1 Receptor Attenuates Early Brain Injury in a Rat Model of Subarachnoid Hemorrhage viathe Suppression of Neuroinflammation through AMPK/TBK1/NF-?B Pathway in Rats Neurotherapeutics 2020-01-01 [PMID: 31486022] (Western Blot)

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-87760

NBP1-87760PEP	NFkB2/NFkB p100 Recombinant Protein Antigen
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-87760

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

