Product Datasheet

ME2 Antibody - BSA Free NBP1-89559

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-89559

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-89559



NBP1-89559

ME2 Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol
Target Molecular Weight	65 kDa
Product Description	
Host	Rabbit
Gene ID	4200
Gene Symbol	MF2

Product Description	
Host	Rabbit
Gene ID	4200
Gene Symbol	ME2
Species	Human
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non-tested species: Mouse (83%), Rat (80%)
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: KVISKPISEHKILFLGAGEAALGIANLIVMSMVENGLSEQEAQKKIWMFDKYGLL VKGRKAKIDSYQEPFTHSAPESIPDTFEDAVNILKPSTIIGVAGAGRLFTPDVIRA MASINERPVIFALSNPTA

Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:20 - 1:50, Immunohistochemistry-Paraffin 1:20 - 1:50
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended.


Images

Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining in human duodenum and liver tissues using anti-ME2 antibody. Corresponding ME2 RNA-seq data are presented for the same tissues.







Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human cerebral cortex, colon, lymph node and testis using Anti-ME2 antibody NBP1-89559 (A) shows similar protein distribution across tissues to independent antibody NBP1-82434 (B).




Western Blot: ME2 Antibody [NBP1-89559] - Analysis using Anti-ME2 antibody NBP1-89559 (A) shows similar pattern to independent antibody NBP1-82434 (B).




Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human colon.




Western Blot: ME2 Antibody [NBP1-89559] - Analysis in human cell line Daudi.




Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human rectum shows strong cytoplasmic positivity in glandular cells.




Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human duodenum shows high expression.




Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human liver shows low expression as expected.



Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human lymph node.



Immunohistochemistry-Paraffin: ME2 Antibody [NBP1-89559] - Staining of human cerebral cortex.



Publications

Brown LJ, Longacre MJ, Hasan NM et al. Chronic reduction of the cytosolic or mitochondrial NAD(P)-malic enzyme does not affect insulin secretion in a rat insulinoma cell line. J Biol Chem. 2009-12-01 [PMID: 19858194]

Zoccarato F, Cavallini L, Alexandre A et al. Succinate is the controller of O2-/H2O2 release at mitochondrial complex I : negative modulation by malate, positive by cyanide. J Bioenerg Biomembr. 2009-08-01 [PMID: 19821037]

MacDonald MJ, Longacre MJ, Kendrick MA et al. Mitochondrial malic enzyme (ME2) in pancreatic islets of the human, rat and mouse and clonal insulinoma cells. Arch Biochem Biophys. 2009-08-01 [PMID: 19691144]

Tsuyuguchi I, Kawasumi H, Takashima T et al. Mycobacterium avium-Mycobacterium intracellular complex-induced suppression of T-cell proliferation in vitro by regulation of monocyte accessory cell activity. Infect Immun. 1990-05-01 [PMID: 1691144]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP1-89559

NBP1-89559PEP ME2 Recombinant Protein Antigen

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-89559

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

