Product Datasheet

Guinea Pig IgG Fc Isotype Control [FITC] NBP1-97257

Unit Size: 1 mg

Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.



Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-97257

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-97257



NBP1-97257

Guinea Pig IgG Fc Isotype Control [FITC]

at -20Č. Avoid freeze-thaw cycles. Preservative 0.01% Sodium Azide Reconstitution Instructions Reconstitute with 1.0 ml deionized water (or equivalent). Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin are against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Applications Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	0 0 1	• •
Concentration LYOPH mg/ml Storage Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles. Preservative 0.01% Sodium Azide Reconstitution Instructions Reconstitute with 1.0 ml deionized water (or equivalent). Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Host Guinea Pig Species Species Guinea Pig Serue Species Guinea Pig Species Specificity/Sensitivity <td< th=""><th>Product Information</th><th></th></td<>	Product Information	
Storage Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles. Preservative 0.01% Sodium Azide Reconstitution Instructions Reconstitute with 1.0 ml deionized water (or equivalent). Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Description Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Applications Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence	Unit Size	1 mg
at -20Č. Avoid freeze-thaw cycles. Preservative 0.01% Sodium Azide Reconstitution Instructions Reconstitute with 1.0 ml deionized water (or equivalent). Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin are against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Applications Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Concentration	LYOPH mg/ml
Reconstitution Instructions Reconstitute with 1.0 ml deionized water (or equivalent). Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Host Guinea Pig Species Guinea Pig Species Guinea Pig Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Applications Fluorophore-linked immunosorbent assay, and fluorescent microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence	Storage	Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.
Isotype IgG Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig Serum Host Guinea Pig Species Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence	Preservative	0.01% Sodium Azide
Conjugate FITC Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig Species Guinea Pig Species Guinea Pig Species Guinea Pig Species Guinea Pig Application Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence	Reconstitution Instructions	Reconstitute with 1.0 ml deionized water (or equivalent).
Purity Multi-step Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig Serum Host Guinea Pig Species Guinea Pig Specificity/Sensitivity No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence	Isotype	IgG
Buffer Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig IgG F(ab')2 or anti- Papain. Host Guinea Pig Species Guinea Pig Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescente microscopy, fluorescence	Conjugate	FITC
10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Product Description Description Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig Serum Host Guinea Pig Specificity/Sensitivity No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Purity	Multi-step
DescriptionStore vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use.This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig SerumHostGuinea PigSpeciesGuinea PigSpecificity/SensitivityNo reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain.Product Application DetailsFluorophore-linked immunosorbent assay, Immunocytochemistry/ ImmunofluorescenceApplication NotesThis product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescence microscopy, fluorescence	Buffer	Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use.This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig SerumHostGuinea PigSpeciesGuinea PigSpecificity/SensitivityNo reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain.Product Application DetailsFluorophore-linked immunosorbent assay, Immunocytochemistry/ ImmunofluorescenceRecommended DilutionsImmunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assayApplication NotesThis product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Product Description	
Species Guinea Pig Specificity/Sensitivity No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Description	freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use. This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Guinea Pig IgG,
Specificity/Sensitivity No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain. Product Application Details Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Host	Guinea Pig
Product Application Details Applications Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Species	Guinea Pig
Applications Fluorophore-linked immunosorbent assay, Immunocytochemistry/ Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Specificity/Sensitivity	No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti- Papain.
Immunofluorescence Recommended Dilutions Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Product Application Details	
Application Notes This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Applications	
based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various	Recommended Dilutions	Immunocytochemistry/ Immunofluorescence, Fluorophore-linked immunosorbent assay
	Application Notes	based plate assays (FLISA) and fluorescent western blotting. This product is also

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-97257

NB7446

Goat anti-Human IgG Fc Secondary Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Isotype Controls are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-97257

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

