

Product Datasheet

SRp55 Antibody - BSA Free

NBP2-04142

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 3 Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-04142

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-04142



NBP2-04142

SRp55 Antibody - BSA Free


Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)

Product Description	
Description	Novus Biologicals Rabbit SRp55 Antibody - BSA Free (NBP2-04142) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-SRp55 Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	6431
Gene Symbol	SRSF6
Species	Human, Mouse
Immunogen	The immunogen recognized by this antibody maps to this antibody using the numbering given in entry NP_006266.2 (GeneID 6431).


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500 - 1:2000
Application Notes	Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections. SRp55 antibody validated for ICC/IF from a verified customer review.

Images


Western Blot: SRp55 Antibody [NBP2-04142] - CEM-T4 and HeLa whole cell lysates. Image from verified customer review.




Immunocytochemistry/Immunofluorescence: SRp55 Antibody [NBP2-04142] - Staining in human fibroblasts. 1X PBS/5% normal serum/0.3% Triton™ X-100. Diluted 1:200 in 1X PBS/1% BSA/0.3% Triton™ X-100, 16 hours at 4C incubation. 2 hours (25C) secondary (1.5:1000 dilution). Image submitted by a verified customer review.




Immunohistochemistry-Paraffin: SRp55 Antibody [NBP2-04142] - Sample: FFPE section of mouse teratoma. Antibody: Affinity purified rabbit anti- SRp55 used at a dilution of 1:1,000 (1ug/ml). Detection: DAB. Counterstain: Hematoxylin (blue).




Western Blot: SRp55 Antibody [NBP2-04142] - analysis of SRp55 in human lymphocytes using anti-SRp55 antibody. Each lane loaded with 14ug of whole cell lysate with 3 different human B-cell lines. Image from verified customer review.



Immunohistochemistry-Paraffin: SRp55 Antibody [NBP2-04142] - Sample: FFPE section of human prostate carcinoma. Antibody: Affinity purified rabbit anti- SRp55 used at a dilution of 1:1,000 (1ug/ml). Detection: DAB. Counterstain: Hematoxylin (blue).



Immunoprecipitation: SRp55 Antibody [NBP2-04142] - Samples: Whole cell lysate from HeLa (15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded), 293T (T; 50 ug), Jurkat (J; 50 ug) and mouse NIH3T3 (M; 50 ug) cells. Antibodies: Affinity purified rabbit anti-SRp55 antibody NBP2-04142 used for WB at 0.1 ug/ml (A) and 1 ug/ml (B) and used for IP at 6 ug/mg lysate. SRp55 was also immunoprecipitated by rabbit anti-SRp55 antibody, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 30 seconds (B).



Publications

Dahal S, Cheng R, Cheung P et al. The Thiazole-5-Carboxamide GPS491 Inhibits HIV-1, Adenovirus, and Coronavirus Replication by Altering RNA Processing/Accumulation *Viruses* 2021-12-30 [PMID: 35062264] (WB, Human)

Holm LL, Doktor TK, Hansen MB et al. Vulnerable exons, like ACADM exon 5, are highly dependent on maintaining a correct balance between splicing enhancers and silencers *Human mutation* 2021-12-19 [PMID: 34923709] (WB)

Wong RW, Balachandran A, Cheung PK et al. An activator of G protein-coupled receptor and MEK1/2-ERK1/2 signaling inhibits HIV-1 replication by altering viral RNA processing *PLoS Pathog.* 2020-02-01 [PMID: 32069328] (WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-04142

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-04142

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

