### **Product Datasheet**

# PKC beta Overexpression Lysate NBP2-04488

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-04488

Updated 11/9/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-04488



#### NBP2-04488

PKC beta Overexpression Lysate

Product Information		
Unit Size	0.1 mg	
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.	
Storage	Store at -80C. Avoid freeze-thaw cycles.	
Buffer	RIPA buffer	
Target Molecular Weight	76.7 kDa	
Product Description		
Description	PKC beta Transient Overexpression Lysate	
	Expression Host: HEK293T	
	Plasmid: RC220599	
	Accession#: NM_212535	
	Protein Tag: C-MYC/DDK	
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.	
Gene ID	5579	
Gene Symbol	PRKCB	
Species	Human	
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.	
Lysate Type	Overexpression	
Product Application Details		
Applications	Western Blot	
Recommended Dilutions	Western Blot	



for 20-	vial of cell lysate contains 100 ug of total protein which should be sufficient
recom	50 reactions. Depending on over-expression level, antibody affinity and on system, some lysates can go as low as 0.1 ug per load. We mend starting with 5ug of cell lysate. Add an equal amount of cell lysate (SDS Sample buffer and boil the SDS samples for 10 minutes before g.

Western Blot: PKC beta Overexpression Lysate (Adult Normal) [NBP2-04488] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PKC beta.

170	-	
130	-	
100	-	
70	-	-
55		-
40	-	
35	-	
25	-	
15	-	
10	_	

www.novusbio.com



technical@novusbio.com



#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-04488

NBP2-38579PEP	PKC beta Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
NBP2-19846	PKC beta Antibody
DVE00	VEGF [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-04488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

