## **Product Datasheet**

### alpha-Taxilin Overexpression Lysate NBP2-05052

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-05052

Updated 11/9/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-05052



#### NBP2-05052

alpha-Taxilin Overexpression Lysate

| Product Information          |  |
|------------------------------|--|
| Unit Size                    | 0.1 mg   |
| Concentration                | The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.   |
| Storage                      | Store at -80C. Avoid freeze-thaw cycles.   |
| Buffer                       | RIPA buffer  |
| Target Molecular Weight      | 61.7 kDa   |
| Product Description          |  |
| Description                  | alpha-Taxilin Transient Overexpression Lysate  |
|                              | Expression Host: HEK293T   |
|                              | Plasmid: RC210301  |
|                              | Accession#: NM_175852  |
|                              | Protein Tag: C-MYC/DDK   |
|                              | You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.  |
| Gene ID                      | 200081   |
| Gene Symbol                  | TXLNA  |
| Species                      | Human  |
| Notes                        | HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid<br>polymer transfection reagent specially designed and manufactured for large<br>volume DNA transfection. Transfected cells were cultured for 48hrs before<br>collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,<br>150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM<br>PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein<br>concentration was measured by BCA protein assay kit. |
| Lysate Type                  | Overexpression   |
| Product Application Details  |  |
| Applications                 | Western Blot   |
| <b>Recommended Dilutions</b> | Western Blot   |
|                              |  |



| fi |  |
|----|--|
| a  | Each vial of cell lysate contains 100ug of total protein which should be sufficient<br>for 20-50 reactions. Depending on over-expression level, antibody affinity and<br>detection system, some lysates can go as low as 0.1 ug per load. We<br>recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate<br>and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before<br>oading. |

#### Images

| Western Blot: alpha-Taxilin Overexpression Lysate [NBP2-05052] - Left-<br>Empty vector transfected control cell lysate (HEK293 cell lysate); Right -<br>Over-expression Lysate for Alpha Taxilin. | 158-<br>106-<br>79-<br>48-<br>35-<br>23- |
|---|--|
|---|--|





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-05052

| NBP1-91662PEP | alpha-Taxilin Recombinant Protein Antigen |
|---------------|---|
| 210-TA-005    | TNF-alpha [Unconjugated]                  |
| NBP1-91662    | alpha-Taxilin Antibody                    |
| 202-IL-010    | IL-2 [Unconjugated]                       |

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-05052

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

