### **Product Datasheet**

# PACE4/PCSK6 Overexpression Lysate NBP2-05676

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-05676

Updated 11/9/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-05676



#### NBP2-05676

PACE4/PCSK6 Overexpression Lysate

| Unit Size 0.1 mg   Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.   Storage Store at -80C. Avoid freeze-thaw cycles.   Buffer RIPA buffer   Target Molecular Weight 47 kDa   Product Description PACE4/PCSK6 Transient Overexpression Lysate   Expression Host: HEK293T Plasmid: RC217415   Accession#: NM_138322 Accession#: NM_138322  |
|--|
| overexpression lysates. Please contact technical support for more information.StorageStore at -80C. Avoid freeze-thaw cycles.BufferRIPA bufferTarget Molecular Weight47 kDaProduct DescriptionPACE4/PCSK6 Transient Overexpression LysateDescriptionExpression Host: HEK293TPlasmid: RC217415<br>Accession#: NM_138322   |
| Buffer RIPA buffer   Target Molecular Weight 47 kDa   Product Description PACE4/PCSK6 Transient Overexpression Lysate   Description Expression Host: HEK293T   Plasmid: RC217415 Accession#: NM_138322   |
| Target Molecular Weight 47 kDa   Product Description PACE4/PCSK6 Transient Overexpression Lysate   Description Expression Host: HEK293T   Plasmid: RC217415 Accession#: NM_138322  |
| Product Description   Description   PACE4/PCSK6 Transient Overexpression Lysate   Expression Host: HEK293T   Plasmid: RC217415   Accession#: NM_138322   |
| Description PACE4/PCSK6 Transient Overexpression Lysate   Expression Host: HEK293T Expression Host: HEK293T   Plasmid: RC217415 Accession#: NM_138322  |
| PACE4/PCSK6 Transient Overexpression Lysate   Expression Host: HEK293T   Plasmid: RC217415   Accession#: NM_138322   |
| Plasmid: RC217415<br>Accession#: NM_138322   |
| Accession#: NM_138322  |
|  |
|  |
| Protein Tag: C-MYC/DDK   |
| You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.  |
| <b>Gene ID</b> 5046  |
| Gene Symbol PCSK6  |
| Species Human  |
| Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. |
| Lysate Type Overexpression   |
| Product Application Details  |
| Applications Western Blot  |
| Recommended Dilutions Western Blot   |



| Application Notes | This product is intended for use as a positive control in Western Blot.<br>Overexpression of the target protein was confirmed using an antibody to DDK<br>(FLAG) epitope tag ( <u>NBP1-71705</u> ) present on the protein construct.  |
|-------------------|---|
|                   | Each vial of cell lysate contains 100ug of total protein which should be sufficient<br>for 20-50 reactions. Depending on over-expression level, antibody affinity and<br>detection system, some lysates can go as low as 0.1 ug per load. We<br>recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate<br>and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before<br>loading. |

#### Images

Western Blot: PACE4 Overexpression Lysate (Adult Normal) [NBP2-05676] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PACE4.

| 170 | _ |   |  |  |
|-----|---|---|--|--|
| 130 | - |   |  |  |
| 100 | _ |   |  |  |
| 70  | - | - |  |  |
| 55  | - |   |  |  |
| 40  | _ |   |  |  |
| 35  | - |   |  |  |
| 25  | - |   |  |  |
| 15  | _ |   |  |  |
| 10  | _ |   |  |  |
|     |   |   |  |  |

www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-05676

| H00005046-Q01-10ug | Recombinant Human PACE4/PCSK6 GST (N-Term) Protein |
|--------------------|--|
| 314-BP-010         | BMP-4 [Unconjugated]                               |
| NBP1-87354         | PACE4/PCSK6 Antibody                               |
| DPC900             | Fibroblast Activation Protein alpha/FAP [HRP]      |

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-05676

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

