Product Datasheet

AUTS2 Overexpression Lysate NBP2-09437

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-09437

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-09437



NBP2-09437

AUTS2 Overexpression Lysate

Unit Size 0.1 mg Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 136.2 kDa Product Description AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate
overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 136.2 kDa Product Description AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
Buffer RIPA buffer Target Molecular Weight 136.2 kDa Product Description AUTS2 Transient Overexpression Lysate Description Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
Target Molecular Weight 136.2 kDa Product Description Description AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
Product Description Description AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
Description AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
AUTS2 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC226365 Accession#: NM_001127231 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control
contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
Gene ID 26053
Gene Symbol AUTS2
Species Human
Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.
Lysate Type Overexpression
Product Application Details
Applications Western Blot
••



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Ы

Western Blot: AUTS2 Overexpression Lysate (Adult Normal) [NBP2-09437] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for AUTS2.

170	-	-
130	-	
95	-	
72	-	110
55	-	
43	-	1
34	-	
26	-	
17	-	
10	_	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-09437

H00026053-Q01-10ug	Recombinant Human AUTS2 GST (N-Term) Protein
AF2377	S100A10 Antibody [Unconjugated]
NBP1-84717	AUTS2 Antibody
NB300-106-100ul	NMDAR2B Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-09437

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

