Product Datasheet

Adenylate Cyclase 3 Overexpression Lysate NBP2-10559

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-10559

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-10559



ermined for ore information.			
Product Description			
n Lysate			
negative control of cell lysate e Buffer consists omophenol blue,			

	of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
Gene ID	109
Gene Symbol	ADCY3

Species	Human
---------	-------

Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid
	polymer transfection reagent specially designed and manufactured for large
	volume DNA transfection. Transfected cells were cultured for 48hrs before
	collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,
	150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM
	PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein

concentration was measured by BCA protein assay kit. Lysate Type Overexpression

Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot




Application Notes

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: Adenylate Cyclase 3 Overexpression Lysate (Adult Normal) [NBP2-10559] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Adenylate Cyclase 3.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-10559

H00000109-Q01-10ug Recombinant Human Adenylate Cyclase 3 GST (N-Term) Protein

AF566 Neuropilin-1 Antibody [Unconjugated]

NBP1-92683 Adenylate Cyclase 3 Antibody

MAB1417 Insulin Antibody (182410) [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-10559

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

