# **Product Datasheet**

# Activin RIIA Overexpression Lysate NBP2-10986

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-10986

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-10986



#### NBP2-10986

Gene ID

**Notes** 

**Gene Symbol** 

Product Information Unit Size 0.1 mg Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more informati Storage Store at -80C. Avoid freeze-thaw cycles.  Buffer RIPA buffer Target Molecular Weight 56 kDa	Activin RIIA Overexpression Lysate		
Concentration  The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information of the protein of interest cannot be determined for overexpression lysates.  Buffer RIPA buffer S6 kDa	duct Information		
overexpression lysates. Please contact technical support for more information Storage  Storage Store at -80C. Avoid freeze-thaw cycles.  Buffer RIPA buffer  Target Molecular Weight 56 kDa	Size		
Buffer RIPA buffer  Target Molecular Weight 56 kDa	centration		
Target Molecular Weight 56 kDa	age		
	er		
	jet Molecular Weight		
Product Description			
Description Activin RIIA Transient Overexpression Lysate	cription		

Expression Host: HEK293T Plasmid: RC210227

Accession#: NM\_001616

Protein Tag: C-MYC/DDK

You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.

92

ACVR2A

Species Human

> HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.

Lysate Type Overexpression

<b>Product Application Details</b>	
Applications	Western Blot
Recommended Dilutions	Western Blot




#### **Application Notes**

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

#### **Images**

Western Blot: Activin RIIA Overexpression Lysate [NBP2-10986] - Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Activin Receptor Type 2A.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-10986**

NBP1-91647PEP Activin RIIA Recombinant Protein Antigen

338-AC-010 Activin A [Unconjugated]

AF340 Activin RIIA Antibody [Unconjugated]

314-BP-010 BMP-4 [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-10986

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

