## **Product Datasheet**

### HLRC1 Overexpression Lysate NBP2-11353

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-11353

Updated 11/9/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-11353



#### NBP2-11353

HLRC1 Overexpression Lysate

Product Information							
Unit Size	0.1 mg						
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.						
Storage	Store at -80C. Avoid freeze-thaw cycles.						
Buffer	RIPA buffer						
Target Molecular Weight	32.7 kDa						
Product Description							
Description	HLRC1 Transient Overexpression Lysate						
	Expression Host: HEK293T						
	Plasmid: RC226938						
	Accession#: NM_001145165						
	Protein Tag: C-MYC/DDK						
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.						
Gene ID	83475						
Gene Symbol	DOHH						
Species	Human						
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.						
Lysate Type	Overexpression						
Product Application Details							
Applications	Western Blot						
Recommended Dilutions	Western Blot						

www.novusbio.com



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag ( <u>NBP1-71705</u> ) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

#### Images

Western Blot: HLRC1 Overexpression Lysate (Adult Normal) [NBP2-11353] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for HLRC1.

170	_		
130	—		
100	_		
70	_		
55	_		
40	_		
35	-	1	
25			
15	-		
10	_		





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-11353

H00083475-P01-10ug	Recombinant Human HLRC1 GST (N-Term) Protein
NB600-326	GST Epitope Tag Antibody - BSA Free
NBP1-91994	HLRC1 Antibody
NBP2-67416	Integrin beta 3/CD61 Antibody (SJ19-09)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-11353

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

