

Product Datasheet

ETS1 associated protein II Antibody Blocking Peptide NBP2-11491PEP

Unit Size: 0.25 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-11491PEP

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-11491PEP



NBP2-11491PEP

ETS1 associated protein II Antibody Blocking Peptide


Product Information	
Unit Size	0.25 mg
Concentration	2.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Purity	>85%, by HPLC
Buffer	Sodium Bicarbonate buffer, pH 8.00 containing up to 10% DMSO

Product Description	
Description	Synthetic peptide corresponding to amino acids 330-360 of human Tyrosyl-DNA phosphodiesterase protein. NCBI Accession #: NP_057698.2 Source: <i>Synthetic</i>
Gene ID	51567
Gene Symbol	TDP2
Species	Human

Product Application Details	
Applications	Antibody Competition
Recommended Dilutions	Antibody Competition
Application Notes	This peptide is useful as a blocking peptide for NBP2-11491. For further blocking peptide related protocol, click here .

Images

Western Blot: ETS1 associated protein II Blocking Peptide [NBP2-11491PEP] - Western Blot of ETS1 Associated protein II antibody with ETS1 associated protein II Blocking peptide in DiluObuffer.



Procedures

Antibody Competition Protocol for ETS1 associated protein II Protein (NBP2-11491PEP)

Antibody Competition Protocol for ETS1 associated protein II Protein (NBP2-11491PEP):

In this assay the antigen binding sites (Fab) of a particular antibody is allowed to bind to homologous antigenic peptide in several hundred molar excess ratio of immunoglobulin to peptide. Since the antigenic peptide has higher affinity for Fab site than the full length native proteins, the antigen binding site is blocked thus antibodies are unable to bind to the specific sites on either native, partial denatured or completely denature antigen on sections, in solution or on membrane blots.

Before starting, standardize the conditions for Western blot, immunoprecipitation or immunohistochemistry using the relevant antibody. Conditions include: volume of antibody used; dilution factor, final volume, incubation time, washing conditions etc. Once conditions have been standardized, repeat the same protocol in duplicate using blocked and unblocked antibody.

Begin the competition assay by preparing the blocked antibody-peptide solution:

1. Take the same volume of antibody as previously standardized. Let us assume 10 uL in 15 mL.
2. Add approximately 1:200 moles of excess peptide (peptides are generally 2200 dalton). This will be equivalent of approximately 60-70ul of peptide solution (original peptide concentration is 2.5mg/ml).
3. Mix 10 uL of antibody with 60-70 uL of peptide and make up the volume to 200 uL using DiluOBuffer.

Compare the above blocked antibody-peptide solution with control antibody:

1. Take same amount of antibody as used in the blocked antibody-peptide solution and add 60-70 ul of PBS and make up the volume to 200 ul using DiluOBuffer.
2. Once both blocked antibody-peptide solution and the control antibody solution have been made, incubate both solutions at 4 degrees C overnight on a rotating mixer. Centrifuge both tubes at 12,000-14000 xg for 1-2 minute at 4 degrees C. A small amount of precipitate may accumulate in the blocked antibody-peptide solution due to antibody-antigen complex formation. If this occurs, carefully remove the precipitate and use only the supernatant.
3. Dilute both solutions in 1X DiluOBuffer to the final dilution volume of 15 mL. The antibody-peptide solution will not give any labeling or will have significantly reduced labeling while the Control Antibody solution should provide standard results.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-11491PEP

NBP2-22913	Recombinant Human ETS1 associated protein II His Protein
210-TA-005	TNF-alpha [Unconjugated]
NBP2-50071-0.1mg	ETS1 associated protein II Antibody (TDP2/1258) - Azide and BSA Free
NB100-116	APE Antibody (13B8E5C2) - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Peptides and proteins are guaranteed for 3 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-11491PEP

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

