Product Datasheet

SLC34A1 Antibody - BSA Free NBP2-13328

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 10

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-13328

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-13328




NBP2-13328

| SLC34A1 Antibody - BSA Free | |
|------------------------------------|--|
| Product Information | |
| Unit Size | 0.1 ml |
| Concentration | Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Clonality | Polyclonal |
| Preservative | 0.02% Sodium Azide |
| Isotype | IgG |
| Purity | Immunogen affinity purified |
| Buffer | PBS (pH 7.2) and 40% Glycerol |
| Product Description | |
| Description | Novus Biologicals Rabbit SLC34A1 Antibody - BSA Free (NBP2-13328) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-SLC34A1 Antibody: Cited in 10 publications. All Novus Biologicals antibodies are covered by our 100% guarantee. |
| Host | Rabbit |
| Gene ID | 6569 |
| Gene Symbol | SLC34A1 |
| Species | Human, Mouse, Rat |
| Reactivity Notes | Mouse reactivity reported in scientific literature (PMID: 26751835), Rat reactivity reported in scientific literature (PMID: 25790436). |
| Immunogen | This antibody was developed against a recombinant protein corresponding to the amino acids: PLPVPGGHVMRGTAFAYVPSPQVLHRIPGTSAYAFPSLGPVALAEHTCPCGEV LERHEPLPAKLALEEEQKPESRLVPKLRQA |
| Product Application Details | |
| Applications | Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry |
| Recommended Dilutions | Western Blot Reported in (PMID: 37746883), Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:200 |
| Application Notes | For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF, Fixation/Permeabilization: PFA/Triton X-100 |
| | |



Images


Immunocytochemistry/Immunofluorescence: SLC34A1 Antibody [NBP2-13328] - Staining of human cell line HEL shows localization to nuclear speckles, plasma membrane & mitotic spindle. Antibody staining is shown in green.




Immunohistochemistry-Paraffin: SLC34A1 Antibody [NBP2-13328] - Staining of human stomach shows no positivity in glandular cells as expected.



Immunohistochemistry: SLC34A1 Antibody [NBP2-13328] - Immunohistochemical analysis of the apical (closed arrows) and intracellular (open arrows) NaPi-2a (SLC34A1) expression in the kidney. Image collected and cropped by CiteAb from the following publication (https://dx.plos.org/10.1371/journal.pone.0116590) licensed under a CC-BY license.



Immunohistochemistry-Paraffin: SLC34A1 Antibody [NBP2-13328] - Staining of human kidney shows moderate cytoplasmic and luminal membranous positivity in cells in tubules.



Page 3 of 5 v.20.1 Updated 9/9/2025 Immunohistochemistry-Paraffin: SLC34A1 Antibody [NBP2-13328] -Staining of human placenta shows no positivity in trophoblastic cells as expected. Immunohistochemistry-Paraffin: SLC34A1 Antibody [NBP2-13328] -Staining of human prostate shows no positivity in glandular cells as expected. Immunohistochemical analysis of the apical (closed arrows) and Vehicle + 0.4 g P intracellular (open arrows) NaPi-2a expression in the kidney. Staining of human cell line HEL shows localization to nuclear speckles, plasma membrane & mitotic spindle.



Publications

Feger M, Alber J, Strotmann J et al. Short-term fasting of mice elevates circulating fibroblast growth factor 23 (FGF23) Acta physiologica (Oxford, England) 2023-09-25 [PMID: 37746883] (WB, Rat)

Huang B, Zeng Z, Li H et al. Modeling kidney development, disease, and plasticity with clonal expandable nephron progenitor cells and nephron organoids bioRxiv 2023-05-26 [PMID: 37293038] (Immunocytochemistry/Immunofluorescence, Human)

Shin S, Boadi EA, Bandyopadhyay BC Ablation of TRPC3 compromises bicarbonate and phosphate transporter activity in mice proximal tubular cells Clinical and experimental pharmacology & physiology 2022-11-26 [PMID: 36433745]

Balzer MS, Doke T, Yang YW et al. Single-cell analysis highlights differences in druggable pathways underlying adaptive or fibrotic kidney regeneration Nature communications 2022-07-11 [PMID: 35821371] (IHC-P, Mouse)

Maruyama H, Taguchi A et al. Low bone mineral density due to secondary hyperparathyroidism in the GlatmTg(CAG-A4GALT) mouse model of Fabry disease. FASEB Bioadv 2020-01-06 [PMID: 32617522] (IF/IHC, WB, Mouse)

Ter Braake AD, Smit AE, Bos C, et al. Magnesium prevents vascular calcification in Klotho deficiency Kidney Int. 2019-11-02 [PMID: 31866113] (WB, Mouse)

Mace ML, Gravesen E, Nordholm A et al. Kidney fibroblast growth factor 23 does not contribute to elevation of its circulating levels in uremia. Kidney Int. 2017-03-21 [PMID: 28341272] (WB, Rat)

Suzuki T, Seki S, Hiramoto K et al. Hyperactivation of Nrf2 in early tubular development induces nephrogenic diabetes insipidus. Nat Commun. 2017-02-24 [PMID: 28233855]

Liu ES, Martins JS, Raimann A et al. 1,25-Dihydroxyvitamin D Alone Improves Skeletal Growth, Microarchitecture and Strength in a Murine Model of XLH, Despite Enhanced FGF23 Expression. J. Bone Miner. Res. 2016-01-11 [PMID: 26751835] (IHC-P, Mouse)

Robijn S, Vervaet BA, D'Haese PC, Verhulst A. Evaluation of intestinal phosphate binding to improve the safety profile of oral sodium phosphate bowel cleansing. PLoS ONE. 2015-03-20 [PMID: 25790436] (IHC-P, Rat)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-13328

NBP2-13328PEP SLC34A1 Recombinant Protein Antigen

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-13328

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

