

Product Datasheet

Cytokeratin 19 Antibody (A53-B/A2.26)

NBP2-15186-0.1mg

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-15186

Updated 4/13/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-15186



NBP2-15186-0.1mg

Cytokeratin 19 Antibody (A53-B/A2.26)

Product Information	
Unit Size	0.1 mg
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	A53-B/A2.26
Preservative	0.05% Sodium Azide
Isotype	IgG2a Lambda
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Target Molecular Weight	40 kDa

Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-33134) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	3880
Gene Symbol	KRT19
Species	Human, Ferret (Negative), Rat (Negative)
Reactivity Notes	Does not react with Rat or Ferret.
Marker	Pancreatic Stem Cell Marker
Specificity/Sensitivity	This monoclonal antibody reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, although 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.
Immunogen	Human breast cancer MCF-7 cells

Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1-2 ug/ml, Simple Western 10 ug/ml, Flow Cytometry 0.5-1 ug/million cells, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Immunohistochemistry-Frozen 0.5-1ug/ml




Application Notes


Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images


Western Blot: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Western Blot Analysis of human Prostate Cancer PC-3 cell lysate Cytokeratin 19 Antibody (A53-B/A2.26).




Immunocytochemistry/Immunofluorescence: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Immunofluorescence Analysis of MCF-7 cells labeling Cytokeratin 19 with Cytokeratin 19 Antibody (A53-B/A2.26) followed by Goat anti-Mouse IgG-CF488 (Cyan). The nuclear counterstain is Red Dot (Red)




Immunohistochemistry-Frozen: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Breast cancer tissue stained for Cytokeratin 19 (red) and counter stained with DAPI (blue). IHC-Fr image submitted by a verified customer review. Image using the Janelia Fluor 549, Azide and BSA free format of this antibody.




Western Blot: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - analysis of Keratin-19 in 1) HepG2 and 2) MCF7 lysate using of Keratin-19 antibody at 1 ug/ml. goat anti-mouse Ig HRP secondary antibody and PicoTect ECL substrate solution were used for this test.




Western Blot: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Western blot 1) Hep-G2 and 2) MCF-7 lysate using of Cytokeratin 19 Antibody (A53-B/A2.26).



Immunocytochemistry/Immunofluorescence: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - HepG2 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton-X100. The cells were incubated with anti-Cytokeratin 19 (A53-B/A2.26) conjugated to Alexa Fluor 488 [NBP2-33134AF488] at 10ug/ml for 1 hour at room temperature. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective. Image from the Alexa Fluor 488 version of this antibody.




Immunohistochemistry-Paraffin: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Formalin-paraffin colon carcinoma stained with Cytokeratin 19 Ab (A53-B/A2.26).



Simple Western: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Simple Western lane view shows a specific band for Cytokeratin 19 in 0.2 mg/ml of HeLa lysate(s). This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Simple Western: Cytokeratin 19 Antibody (A53-B/A2.26) [NBP2-15186] - Electropherogram image of the corresponding Simple Western lane. Cytokeratin 19 antibody was used at 10 ug/ml dilution of HeLa lysates(s) respectively.



Publications

Wu Q, Hu Y, Ma Q et al. Comprehensive Analysis of RAPGEF2 for Predicting Prognosis and Immunotherapy Response in Patients with Hepatocellular Carcinoma Journal of Oncology 2022-04-26 [PMID: 35518785] (FLOW)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-15186-0.1mg

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-78278PEP	Cytokeratin 19 Antibody Blocking Peptide
1129-ER-050	ErbB2/Her2 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-15186

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

