

Product Datasheet

Complement Component C9 Antibody - BSA Free NBP2-15952

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-15952

Updated 2/23/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-15952




NBP2-15952

Complement Component C9 Antibody - BSA Free


| Product Information | |
|-----------------------------|---|
| Unit Size | 0.1 ml |
| Concentration | Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services. |
| Storage | Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles. |
| Clonality | Polyclonal |
| Preservative | 0.025% Proclin 300 |
| Isotype | IgG |
| Purity | Antigen Affinity-purified |
| Buffer | PBS, 20% Glycerol |
| Target Molecular Weight | 63 kDa |
| Product Description | |
| Host | Rabbit |
| Gene ID | 735 |
| Gene Symbol | C9 |
| Species | Human, Mouse, Rat |
| Immunogen | Recombinant protein encompassing a sequence within the center region of human Complement Component C9. The exact sequence is proprietary. |
| Product Application Details | |
| Applications | Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry |
| Recommended Dilutions | Western Blot 1:500-1:3000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100-1:1000 |

Images


Immunohistochemistry: Complement Component C9 Antibody [NBP2-15952] - A generic staining of the vascular BMs was given by an antibody to the 7S domain of collagen IV $\alpha 3$ (C). Prominent staining for microvascular aneurisms was detected by staining with antibodies to C9 (B, C). The same treatment of vascular BM whole mounts from non-diabetic eyes did not show staining for these proteins (D). Image collected and cropped by CiteAb from the following publication (dx.plos.org/10.1371/journal.pone.0189857) licensed under a CC-BY license.




Western Blot: Complement Component C9 Antibody [NBP2-15952] - Sample (30 ug of whole cell lysate) A: THP-1 7.5% SDS PAGE gel, diluted at 1:500.




Immunocytochemistry/Immunofluorescence: Complement Component C9 Antibody [NBP2-15952] - HepG2 cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: C9 stained by C9 antibody [N2C2-2], Internal diluted at 1:500. Blue: Hoechst 33342 staining.




Western Blot: Complement Component C9 Antibody [NBP2-15952] - Human plasma (30 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with C9 antibody [N2C2-2], Internal diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Western Blot: Complement Component C9 Antibody [NBP2-15952] - Rat plasma (50 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with Complement Component C9 Antibody diluted at 1:1000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Western Blot: Complement Component C9 Antibody [NBP2-15952] - Mouse plasma (50 ug) was separated by 12% SDS-PAGE, and the membrane was blotted with Complement Component 9 antibody diluted at 1:1000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody,




Immunocytochemistry/ Immunofluorescence: Complement Component C9 Antibody [NBP2-15952] - Staining of vascular BM whole mounts with antibodies to proteins detected in the proteome analysis. A generic staining of the vascular BMs was given by an antibody to the 7S domain of collagen IV α 3 (A, C, E, F, G). Prominent staining for microvascular aneurisms was detected by staining with antibodies to C9 (B, C), Fibronectin (FN, E), ApoE (F) & PRELP (G). The same treatment of vascular BM whole mounts from non-diabetic eyes did not show staining for these proteins (D). A norrin-specific staining is shown to be generic for the entire vascular BM whole mounts (H), the signal, however, being less prominent in vascular aneurisms (arrow in H). Staining of vascular BM whole mounts from non-diabetic eyes showed a clearly weaker staining for norrin, when compared to vascular whole mounts from non-diabetic donors. Bar: 25um. Image collected & cropped by CiteAb from the following publication

(<https://dx.plos.org/10.1371/journal.pone.0189857>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: Complement Component C9 Antibody [NBP2-15952] - Staining of vascular BM whole mounts with antibodies to proteins detected in the proteome analysis. A generic staining of the vascular BMs was given by an antibody to the 7S domain of collagen IV α 3 (A, C, E, F, G). Prominent staining for microvascular aneurisms was detected by staining with antibodies to C9 (B, C), Fibronectin (FN, E), ApoE (F) & PRELP (G). The same treatment of vascular BM whole mounts from non-diabetic eyes did not show staining for these proteins (D). A norrin-specific staining is shown to be generic for the entire vascular BM whole mounts (H), the signal, however, being less prominent in vascular aneurisms (arrow in H). Staining of vascular BM whole mounts from non-diabetic eyes showed a clearly weaker staining for norrin, when compared to vascular whole mounts from non-diabetic donors. Bar: 25um. Image collected & cropped by CiteAb from the following publication

(<https://dx.plos.org/10.1371/journal.pone.0189857>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: Complement Component C9 Antibody [NBP2-15952] - Staining of vascular BM whole mounts with antibodies to proteins detected in the proteome analysis. A generic staining of the vascular BMs was given by an antibody to the 7S domain of collagen IV α 3 (A, C, E, F, G). Prominent staining for microvascular aneurisms was detected by staining with antibodies to C9 (B, C), Fibronectin (FN, E), ApoE (F) & PRELP (G). The same treatment of vascular BM whole mounts from non-diabetic eyes did not show staining for these proteins (D). A norrin-specific staining is shown to be generic for the entire vascular BM whole mounts (H), the signal, however, being less prominent in vascular aneurisms (arrow in H). Staining of vascular BM whole mounts from non-diabetic eyes showed a clearly weaker staining for norrin, when compared to vascular whole mounts from non-diabetic donors. Bar: 25um. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0189857>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Halfter W, Moes S, Asgeirsson DO et al. Diabetes-related changes in the protein composition and the biomechanical properties of human retinal vascular basement membranes. PLoS ONE 2017-12-28 [PMID: 29284024] (ICC/IF, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-15952

| | |
|--------------------|--|
| HAF008 | Goat anti-Rabbit IgG Secondary Antibody [HRP] |
| NB7160 | Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP] |
| NBP2-24891 | Rabbit IgG Isotype Control |
| H00000735-P01-10ug | Recombinant Human Complement Component C9 GST (N-Term) Protein |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-15952

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

