# **Product Datasheet**

# DCK Antibody - BSA Free NBP2-16108

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-16108

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-16108



### NBP2-16108

DCK Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS (pH 7), 20% Glycerol
Target Molecular Weight	31 kDa
Product Description	
Description	Novus Biologicals Rabbit DCK Antibody - BSA Free (NBP2-16108) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-DCK

<b>Product Description</b>	
Description	Novus Biologicals Rabbit DCK Antibody - BSA Free (NBP2-16108) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-DCK Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1633
Gene Symbol	DCK
Species	Human, Mouse, Rat
Reactivity Notes	Chicken (82%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human DCK. The exact sequence is proprietary.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:20000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunoprecipitation 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:1000


## **Images**

Western Blot: DCK Antibody [NBP2-16108] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody. HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.







Western Blot: DCK Antibody [NBP2-16108] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody diluted at 1:5000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: DCK Antibody [NBP2-16108] - HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: DCK protein stained by DCK antibody diluted at 1:500. Red: Phalloidin, a cytoskeleton marker, diluted at 1:100. Scale bar = 10 um.




Immunohistochemistry-Paraffin: DCK Antibody [NBP2-16108] - Rat lung. DCK stained by DCK antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Western Blot: DCK Antibody [NBP2-16108] - Various whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody diluted at a dilution of 1:5000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Western Blot: DCK Antibody [NBP2-16108] - A. 30 ug PC-12 whole cell lysate/extract B. 30 ug Rat2 whole cell lysate/extract 12% SDS-PAGE DCK antibody dilution: 1:1000 The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Immunohistochemistry-Paraffin: DCK Antibody [NBP2-16108] - Mouse hind brain. DCK antibody dilution: 1:500. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.



Immunoprecipitation: DCK Antibody [NBP2-16108] - Immunoprecipitation of DCK protein from 293T whole cell extracts using 5 ug of DCK antibody. Western blot analysis was performed using DCK antibody. EasyBlot anti-Rabbit IgG was used as a secondary reagent.



Western Blot: DCK Antibody [NBP2-16108] - Various whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody (NBP2-16108) diluted at 1:3000. The HRPconjugated anti-rabbit IgG antibody was used to detect the primary antibody.



#### **Publications**

Chen YH, Chen YC, Lin CC et al. Synergistic Anticancer Effects of Gemcitabine with Pitavastatin on Pancreatic Cancer Cell Line MIA PaCa-2 in vitro and in vivo Cancer Manag Res 2020-06-17 [PMID: 32606957]





## **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-16108**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-49305-0.1mg Recombinant Human DCK His Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-16108

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

