

Product Datasheet

N-Cadherin Antibody NBP2-19459

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-19459

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-19459



NBP2-19459

N-Cadherin Antibody


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS (pH 7), 20% Glycerol, 1% BSA
Target Molecular Weight	100 kDa

Product Description	
Host	Rabbit
Gene ID	1000
Gene Symbol	CDH2
Species	Human, Mouse, Rat
Reactivity Notes	Xenopus laevis (82%).
Marker	Mesenchymal Cells Marker
Immunogen	Recombinant protein encompassing a sequence within the center region of human N-Cadherin. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockout Validated
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Knockout Validated

Images


Knockout Validated: N-Cadherin Antibody [NBP2-19459] - Wild-type (WT) and N-Cadherin knockout (KO) 293T cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with N-Cadherin antibody diluted at 1:500. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: N-Cadherin Antibody [NBP2-19459] - N-Cadherin antibody detects N-Cadherin protein at cell membrane by immunofluorescent analysis. Sample: NT2D1 cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: N-Cadherin stained by N-Cadherin antibody diluted at 1:500. Blue: Hoechst 33342 staining.




Western Blot: N-Cadherin Antibody [NBP2-19459] - Various tissue extracts (50 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with N-Cadherin antibody [N1N3] diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Immunohistochemistry-Paraffin: N-Cadherin Antibody [NBP2-19459] - Mouse liver. N-Cadherin stained by N-Cadherin antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Western Blot: N-Cadherin Antibody [NBP2-19459] - Rat tissue extract (50 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with N-cadherin antibody (diluted at 1:1000). The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Western Blot: N-Cadherin Antibody [NBP2-19459] - A. 30 ug PC-12 whole cell extract B. 30 ug Rat2 whole cell extract 5% SDS-PAGE N-Cadherin antibody [N1N3] dilution: 1:1000 The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Western Blot: N-Cadherin Antibody [NBP2-19459] - Whole cell extract (30 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with N-Cadherin antibody [N1N3] diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Western Blot: N-Cadherin Antibody [NBP2-19459] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with N-Cadherin antibody [N1N3] diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody



Immunohistochemistry-Paraffin: N-Cadherin Antibody [NBP2-19459] - Human hepatoma. N-Cadherin antibody [N1N3] dilution: 1:500. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.



Immunohistochemistry-Paraffin: N-Cadherin Antibody [NBP2-19459] - Rat liver. N-Cadherin stained by N-Cadherin antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.



Publications

Po JW, Roohullah A, Lynch D et al. Improved ovarian cancer EMT-CTC isolation by immunomagnetic targeting of epithelial EpCAM and mesenchymal N-cadherin. *J Circ Biomark* 2018-06-24 [PMID: 30013673] (ICC/IF, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-19459

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-59927-50ug	Recombinant Human N-Cadherin His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-19459

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

