# **Product Datasheet**

## NDP52 Antibody - BSA Free NBP2-19499

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-19499

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-19499



#### NBP2-19499

NDP52 Antibody - BSA Free

<b>Product Information</b>	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 20% Glycerol
Target Molecular Weight	52 kDa
<b>Product Description</b>	
Description	Novus Biologicals Rabbit NDP52 Antibody - BSA Free (NBP2-19499) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	10241
Gene Symbol	CALCOCO2
Species	Human
Immunogen	Recombinant protein encompassing a sequence within the center region of human NDP52. The exact sequence is proprietary.
<b>Product Application Details</b>	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500-1:10000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunoprecipitation


1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:1000

### **Images**


Western Blot: NDP52 Antibody [NBP2-19499] - NDP52 antibody detects NDP52 protein by western blot analysis.


Un-treated (-) and treated (+, Thapsigargin treatment for 12hrs and 24hrs) HepG2 whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody. The ACTB was used as internal control, shown at the bottom panel.

The ACTB was used as internal control, shown at the bottom panel. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: NDP52 Antibody [NBP2-19499] - NDP52 antibody detects NDP52 protein at autophagosome by immunofluorescent analysis. Sample: Mock and treated HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: NDP52 stained by NDP52 antibody diluted at 1:1000. Red: phalloidin, a cytoskeleton marker, diluted at 1:200. Blue: Fluoroshield with DAPI. Scale bar= 10 um.





Immunohistochemistry-Paraffin: NDP52 Antibody [NBP2-19499] - H441 xenograft, using NDP52 antibody at 1:100 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.




Western Blot: NDP52 Antibody [NBP2-19499] - Sample (30 ug of whole cell lysate) A: Raji 10% SDS PAGE gel, diluted at 1:1000.




Western Blot: NDP52 Antibody [NBP2-19499] - NDP52 antibody detects NDP52 protein by western blot analysis.


Un-treated (-) and treated (+), Thapsigargin treatment for 12hrs and 24hrs. Huh-7 whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody. ACTB was used as internal control, shown at the bottom panel. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Immunoprecipitation: NDP52 Antibody [NBP2-19499] - Jurkat whole cell extracts using 5 ug of NDP52 antibody. Western blot analysis was performed using NDP52 antibody. EasyBlot anti-Rabbit IgG was used as a secondary reagent.



Western Blot: NDP52 Antibody [NBP2-19499] - Untreated (-) and treated (+) HepG2 whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody (NBP2-19499) diluted at 1:2000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP2-19499**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00010241-P01-10ug Recombinant Human NDP52 GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-19499

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

