

Product Datasheet

NGFR/TNFRSF16/p75NTR Antibody NBP2-19669

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-19669

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-19669



NBP2-19669**NGFR/TNFRSF16/p75NTR Antibody**


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS (pH 7), 20% Glycerol, 1% BSA
Target Molecular Weight	45 kDa

Product Description	
Host	Rabbit
Gene ID	4804
Gene Symbol	NGFR
Species	Human, Mouse, Rat
Immunogen	Recombinant protein encompassing a sequence within the center region of human NGFR/TNFRSF16/p75NTR. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:500-1:3000, Flow Cytometry, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Immunohistochemistry-Frozen 1:100-1:1000
Application Notes	Flow-Assay dependent (Membrane permeabilization is required for this application).

Images


Western Blot: NGFR/TNFRSF16/p75NTR Antibody [NBP2-19669] - Untreated (-) and treated (+) PC-3 whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with p75 NGF Receptor / CD271 antibody diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: NGF R/TNFRSF16/p75NTR Antibody [NBP2-19669] - Culture rat E18 primary cortical neuron, DIV 8. Cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: p75 NGF Receptor protein stained by p75 NGF Receptor antibody diluted at 1:250. Red: beta Tubulin 3/ TUJ1, stained by beta Tubulin 3/ TUJ1 antibody [11710] (NBP2-43559) diluted at 1:250. Blue: Fluoroshield with DAPI.




Immunohistochemistry-Frozen: NGF R/TNFRSF16/p75NTR Antibody [NBP2-19669] - Frozen sectioned E13.5 Rat brain. Green: p75 NGF Receptor protein stained by p75 NGF Receptor antibody diluted at 1:250. Red: beta Tubulin 3/ TUJ1, a mature neuron marker, stained by beta Tubulin 3/ TUJ1 antibody [11710] diluted at 1:500. Blue: Fluoroshield with DAPI.




Western Blot: NGF R/TNFRSF16/p75NTR Antibody [NBP2-19669] - Whole cell extracts (30 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with p75 NGF Receptor antibody at a dilution of 1:2000.




Western Blot: NGF R/TNFRSF16/p75NTR Antibody [NBP2-19669] - Various tissue extracts (50 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with p75 NGF Receptor / CD271 antibody diluted at 1:2000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.



Immunohistochemistry-Paraffin: NGF R/TNFRSF16/p75NTR Antibody [NBP2-19669] - Rat brain. p75 NGF Receptor antibody diluted at 1:500.



Western Blot: NGFR/TNFRSF16/p75NTR Antibody [NBP2-19669] - Untreated (-) and treated (+) PC-3 whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with NGFR/TNFRSF16/p75NTR antibody (NBP2-19669) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Publications

Lotti R, Palazzo E, Quadri M et al. Isolation of an "Early" Transit Amplifying Keratinocyte Population in Human Epidermis: A Role for the Low Affinity Neurotrophin Receptor CD271 Stem Cells 2022-12-31 [PMID: 36037263] (Simple Western)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-19669

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-52198-0.05mg	Recombinant Human NGFR/TNFRSF16/p75NTR His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-19669

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

