

Product Datasheet

RUVBL1 Antibody NBP2-20245

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-20245

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-20245



NBP2-20245**RUVBL1 Antibody****Product Information**

Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS (pH 7), 20% Glycerol, 1% BSA
Target Molecular Weight	50 kDa

Product Description


Description	Novus Biologicals Rabbit RUVBL1 Antibody (NBP2-20245) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-RUVBL1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	8607
Gene Symbol	RUVBL1
Species	Human, Mouse, Rat
Reactivity Notes	Xenopus laevis (97%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human RUVBL1. The exact sequence is proprietary.

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Knockdown Validated
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunoprecipitation 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:1000, Knockdown Validated Reported in scientific literature (PMID: 31018511).

Images


Western Blot: RUVBL1 Antibody [NBP2-20245] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with RUVBL1 antibody diluted at 1:2500. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: RUVBL1 Antibody [NBP2-20245] - HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: RUVBL1 protein stained by RUVBL1 antibody diluted at 1:500. Blue: Hoechst 33342 staining. Scale bar = 10 μ m.




Immunohistochemistry-Paraffin: RUVBL1 Antibody [NBP2-20245] - Mouse testis. RUVBL1 antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Immunohistochemistry-Paraffin: RUVBL1 Antibody [NBP2-20245] - Human colon carcinoma, using RUVBL1 antibody at 1:500 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.



Immunohistochemistry-Paraffin: RUVBL1 Antibody [NBP2-20245] - Rat heart. RUVBL1 antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Immunohistochemistry-Paraffin: RUVBL1 Antibody [NBP2-20245] - Mouse colon. RUVBL1 antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Immunohistochemistry-Paraffin: RUVBL1 Antibody [NBP2-20245] - Rat testis. RUVBL1 antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Immunoprecipitation: RUVBL1 Antibody [NBP2-20245] - Immunoprecipitation of RUVBL protein from Jurkat whole cell extracts using 5 ug of RUVBL1 antibody. Western blot analysis was performed using RUVBL1 antibody. EasyBlot anti-Rabbit IgG was used as a secondary reagent.




RUVBL1/2 do not effect EBOV minigenome activity. (A) Schematic diagram of the EBOV minigenome system. The EBOV minigenome system consists of six plasmids: Four support plasmids encode replication complex components NP, L, VP35, & VP30. The EBOV minigenome plasmid encodes a firefly luciferase reporter gene flanked by the leader & trailer sequences of EBOV, & the plasmid that encodes Renilla luciferase is used for normalization. (B) Minigenome activity upon the knockdown of either RUVBL1, RUVBL2, or in combination. Below are protein levels confirmed by immunoblot. HeLa cells were transfected with 80 nM scrambled siRNA, 30 nM siRNA targeting RUVBL1, or 50 nM siRNA targeting RUVBL2. Twenty-four h after siRNA addition, the minigenome components were transfected. Forty-eight h later, minigenome reporter activity was measured. (C) Overexpression of FLAG-RUVBL1 & HA-RUVBL2 in the EBOV minigenome. HeLa cells were left untransfected, or transfected with vector control (VC), or increasing amounts of FLAG-RUVBL1 (125, 250, & 500 ng) or HA-RUVBL2 (125, 250, & 500 ng). Twenty-four h after exogenous transfection, the minigenome components were transfected. Forty-eight h later, minigenome reporter activity was measured. Data represent mean \pm SEM from one representative experiment (n = 3) of at least three experiments (* p < 0.05). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31018511>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: RUVBL1 Antibody [NBP2-20245] - Endogenous RUVBL1 & RUVBL2 colocalize with HA-NP. HeLa cells were transfected with vector control, HA-NP, or HA-VP35. Twenty-four h later, the cells were fixed & processed for immunofluorescence detection of endogenous RUVBL1 or RUVBL2 in the presence of vector control, HA-NP, or HA-VP35. Representative images of (A) endogenous RUVBL1 localization pattern with control vector (top panels), HA-NP (middle panels), or HA-VP35 (bottom panels) & (B) endogenous RUVBL2 localization pattern with control vector (top panels), HA-NP (middle panels), or HA-VP35 (bottom panels) are shown. HA-NP or HA-VP35 (green), RUVBL1/2 (red), & Hoechst 33342 nuclear stain (blue) were visualized by confocal microscopy. Scale bars = 20 μ M. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31018511>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: RUVBL1 Antibody [NBP2-20245] - Various whole cell extracts (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with RUVBL1 antibody (NBP2-20245) diluted at 1:1000.



Publications

Silva B, Pentz R, Figueira AM et al. Identification of RUVBL1 and RUVBL2 as Novel Cellular Interactors of the Ebola Virus Nucleoprotein Viruses 2019-04-23 [PMID: 31018511] (WB, ICC/IF, KD, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-20245

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-50845-0.1mg	Recombinant Human RUVBL1 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-20245

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

