Product Datasheet

Transgelin/TAGLN/SM22 alpha Antibody - BSA Free NBP2-20405

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-20405

Updated 2/23/2025 v.20.1


Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-20405




NBP2-20405


Transgelin/TAGLN/SM22 alpha	Antibody - BSA Free
Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS (pH 7), 20% Glycerol
Target Molecular Weight	23 kDa
Product Description	
Host	Rabbit
Gene ID	6876
Gene Symbol	TAGLN
Species	Human, Mouse, Rat, Bovine
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non-tested species: Chicken (87%), Xenopus laevis (81%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human Transgelin/TAGLN/SM22 alpha. The exact sequence is proprietary.
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:500-1:10000, Immunohistochemistry 1:100, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Immunohistochemistry-Frozen
Images	
_	I/SM22 alpha Antibody [NBP2-20405] - KDa
0 1 (50 (1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	A A



Immunocytochemistry/Immunofluorescence: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: Transgelin protein stained by Transgelin antibody diluted at 1:100. Red: alpha Tubulin, a sytoskeleton marker, stained by alpha Tubulin antibody NBP2-43837) diluted at 1:500. Blue: Hoechst 33342 staining.




Immunohistochemistry: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Coronary artery vascular smooth muscle cells in fixed, frozen adult mouse heart. Primary antibody 1:100 dilution of a 0.2mg/ml solution. This image was submitted via customer Review.



Western Blot: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Western blot analysis of Transgelin and beta-actin (lower panel) Sample (35 ug of whole cell lysate)

A: A-10 cell


B: A-10 cell treated with MG132 NBP2-20405.




Immunohistochemistry-Paraffin: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Paraffin-embedded mouse heart. Transgelin antibody dilution: 1:500.




Immunohistochemistry-Paraffin: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Paraffin-embedded mouse esophagus. Transgelin antibody dilution: 1:500.




Immunohistochemistry-Paraffin: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - 40X image of coronary artery vascular smooth muscle stained for transgelin. Primary Antibody dilution of 1:100. This image was submitted via customer Review.




Western Blot: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Various whole cell extracts (30 ug) were separated by 15% SDS-PAGE, and the membrane was blotted with Transgelin/TAGLN/SM22 alpha antibody (NBP2-20405) diluted at 1:1000. The HRP-conjugated antirabbit IgG antibody was used to detect the primary antibody.



Immunohistochemistry-Paraffin: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Transgelin/TAGLN/SM22 alpha antibody detects Transgelin/TAGLN/SM22 alpha protein at cytoplasm by immunohistochemical analysis.Sample: Paraffin-embedded mouse smooth muscle.Transgelin/TAGLN/SM22 alpha stained by Transgelin/TAGLN/SM22 alpha antibody (NBP2-20405) diluted at 1:500.Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



Western Blot: Transgelin/TAGLN/SM22 alpha Antibody [NBP2-20405] - Various whole cell extracts (30 ug) were separated by 15% SDS-PAGE, and the membrane was blotted with Transgelin antibody diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody. Corresponding RNA expression data for the same cell lines are based on Human Protein Atlas program.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-20405

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-45267-0.1mg Recombinant Human Transgelin/TAGLN/SM22 alpha His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-20405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

