## **Product Datasheet**

# alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) - BSA Free NBP2-22397

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-22397

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-22397



alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) - BSA Free	
Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	S330A-80
Preservative	0.09% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS (pH 7.4), 50% Glycerol
Product Description	
Description	Novus Biologicals Mouse alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) - BSA Free (NBP2-22397) is a monoclonal antibody validated for use in WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	152
Gene Symbol	ADRA2C
Species	Human, Mouse
Specificity/Sensitivity	Detects 50kDa or larger (possibly due to dimerization). Does not cross-react with other adrenergic receptors.
Immunogen	Synthetic peptide amino acids 442- 462 (QDFRRSFKHILFRRRRGFRQ, cytoplasmic C-terminus) of human Alpha-2C adrenergic receptor
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Immunocytochemistry/ Immunofluorescence 1:10-1:2000
Application Notes	1 ug/ml of Alpha-2C Adrenergic Receptor Antibody was sufficient for detection of Alpha2C Adrenergic Receptor in 20 ug of COS cells transiently transfected with HA-tagged Alpha- lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary Antibody.



#### **Images**

Western Blot: alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) [NBP2-22397] - Western Blot analysis of Monkey COS cells transfected with HA-tagged Alpha-2C showing detection of ~50 kDa Alpha-2C Adrenergic Receptor protein using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 (NBP2-22397). Lane 1: Molecular Weight Ladder. Lane 2: Monkey COS cells transfected with HA-tagged Alpha-2C. Load: 15 ug. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody (NBP2-22397) at 1:200 for 16 hours at 4C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~50 kDa.

kDa ww 2


150 ──
100 ──
75 ──
50 ──
37 ──

\*\*Comparison of the comparison of the co


Immunocytochemistry/Immunofluorescence: alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) [NBP2-22397] - Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 (NBP2-22397). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4% PFA for 15 min. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody (NBP2-22397) at 1:100 for overnight at 4C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain; Hoechst (blue) nuclear stain at 1:800, 1.6mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Alpha-2C Adrenergic Receptor Antibody (D) Composite.

A B

Immunocytochemistry/Immunofluorescence: alpha-2C Adrenergic R/ADRA2C Antibody (\$330A-80) [NBP2-22397] - Tissue: Neuroblastoma cell line SK-N-BE. Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X.



Immunocytochemistry/Immunofluorescence: alpha-2C Adrenergic R/ADRA2C Antibody (S330A-80) [NBP2-22397] - Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 (NBP2-22397). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody (NBP2-22397) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) Alpha-2C Adrenergic Receptor Antibody. (D) Composite.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP2-22397**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

NBP2-33845PEP alpha-2C Adrenergic R/ADRA2C Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-22397

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

