

# Product Datasheet

## HSP60 Antibody (4B9/89) - Azide and BSA Free NBP2-22440

Unit Size: 100ug

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-22440](http://www.novusbio.com/NBP2-22440)

Updated 10/23/2024 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-22440](http://www.novusbio.com/reviews/destination/NBP2-22440)



**NBP2-22440**

HSP60 Antibody (4B9/89) - Azide and BSA Free


Product Information	
Unit Size	100ug
Concentration	LYOPH mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4B9/89
Preservative	No Preservative
Reconstitution Instructions	Reconstitute with PBS.
Isotype	IgG2a
Purity	Protein A purified
Buffer	PBS

Product Description	
Host	Mouse
Gene ID	3329
Gene Symbol	HSPD1
Species	Human, Mouse, Rat, Primate
Marker	Mitochondria Marker
Immunogen	Human placental HSP60.


Product Application Details	
Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:100 - 1:1000, ELISA 1:100 - 1:2000, Immunohistochemistry 1:200, Immunocytochemistry/ Immunofluorescence 1:10-1:500, Immunoprecipitation 2 ug, Immunohistochemistry-Paraffin 1:200

**Images**


Western Blot: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of 50ug of the indicated whole cell lysates and 15ul of PageRuler Prestained Protein Ladder.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 (HSP60) (green) in HeLa and NIH3T3 cells. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with a HSP60 monoclonal antibody, at a dilution of 1:50 for at least 1 hour at room temperature, washed with PBS, and incubated with DyLight 488 goat-anti-mouse IgG secondary antibody at a dilution of 1:400 for 30 minutes at room temperature. Nuclei (blue) were stained with Hoechst 33342 dye.




Immunohistochemistry-Paraffin: HSP60 Antibody (4B9/89) [NBP2-22440] - Normal deparaffinized human Tonsil tissue.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 using Anti-Heat Shock Protein 60 Monoclonal Antibody (4B9/89) shows staining in A2058 Cells. Heat Shock Protein 60 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Heat Shock Protein 60 at a dilution of 1:200 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 using Anti-Heat Shock Protein 60 Monoclonal Antibody (4B9/89) shows staining in ATDC5 Cells. Heat Shock Protein 60 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Heat Shock Protein 60 at a dilution of 1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 using Anti-Heat Shock Protein 60 Monoclonal Antibody (4B9/89) shows staining in HeLa Cells. Heat Shock Protein 60 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Heat Shock Protein 60 at a dilution of 1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 using a Hsp60 Monoclonal Antibody in human endothelial cells.




Immunocytochemistry/Immunofluorescence: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 (Hsp60, green) in HeLa cells. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with (left panel) or without (right panel) a Hsp60 monoclonal antibody, at a dilution of 1:50 for at least 1 hour at room temperature, washed with PBS, and incubated with DyLight 488 goat-anti-mouse IgG secondary antibody at a dilution of 1:400 for 30 minutes at room temperature. F-Actin (red) was stained with Dylight 554 phalloidin, and nuclei (blue) were stained with Hoechst 33342 dye.



Immunohistochemistry-Paraffin: HSP60 Antibody (4B9/89) [NBP2-22440] - Cancer biopsies of deparaffinized human Breast carcinoma tissue.



Immunohistochemistry-Paraffin: HSP60 Antibody (4B9/89) [NBP2-22440] - Normal deparaffinized human Kidney tissue.



Immunoprecipitation: HSP60 Antibody (4B9/89) [NBP2-22440] - Analysis of Heat Shock Protein 60 (HSP60) was performed on HeLa cells. Antigen:antibody complexes were formed by incubating 500ug whole cell lysate with 2ug of HSP60 monoclonal antibody overnight on a rocking platform at 4C. The immune complexes were captured on 50ul Protein A/G Plus Agarose, washed extensively, and eluted with Lane Marker Reducing Sample Buffer. Samples were then resolved on a 4-20% Tris-HCl polyacrylamide gel, transferred to a PVDF membrane, and blocked with 5% BSA/TBST for at least 1 hour. The membrane was probed with a HSP60 monoclonal antibody at a dilution of 1:1000 overnight rotating at 4C, washed in TBST, and probed with Clean-Blot IP Detection Reagent (HRP) at a dilution of 1:1000 for at least one hour. Chemiluminescent detection was performed using SuperSignal West Dura.







### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-22440**

---

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-96778	Mouse IgG2a Isotype Control (M2A)
NBP1-77396PEP	HSP60 Antibody Blocking Peptide

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-22440](http://www.novusbio.com/reviews/submit/NBP2-22440)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

