# **Product Datasheet**

# Smad7 Antibody - BSA Free NBP2-24710

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.



#### **Publications: 14**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-24710

Updated 2/23/2025 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-24710



## NBP2-24710

Smad7 Antibody - BSA Free

Product Information			
Unit Size	0.1 mg		
Concentration	1.0 mg/ml		
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.		
Clonality	Polyclonal		
Preservative	0.05% Sodium Azide		
Isotype	IgG		
Purity	Protein G purified		
Buffer	PBS		
Product Description			
Host	Rabbit		
Gene ID	4092		
Gene Symbol	SMAD7		
Species	Human, Mouse, Sheep		
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 24576683).		
Specificity/Sensitivity	This antibody may cross-react with SMAD6 depending on cell or tissue-specific expression.		
Immunogen	This antibody was raised against two peptides made from within amino acids 10 and 50 of human SMAD7. This sequence is 100% or highly conserved between human and multiple species.		
Product Application Details			
Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation		
Recommended Dilutions	Western Blot 2-5 ug/ml, ELISA reported in scientific literature (PMID 32108968), Immunohistochemistry 5-10 ug/ml, Immunoprecipitation, Immunohistochemistry- Paraffin 5 ug/ml		



pSmad3

Smad3

pSmad2

Smad2/3

Smad4

Smad7

a-tubulin

#### Images

Western Blot: Smad7 Antibody [NBP2-24710] - Effect of mTOR inhibitors on TGF-beta activation of mTOR and Smad pathways. Serum-deprived IPF fibroblasts were treated with or without TGF-beta for 15 minutes in the presence or absence of indicated inhibitors followed by Western blot analysis with an anti-phospho-Smad2 or Smad3 antibody. Expression of total Smad-2, 3, 4 and 7 was analyzed by Western blot. Experiment was done on three lines, which are shown in Figure 1; results were similar between the three lines and results from the IPF fibroblasts isolated from surgical lung biopsy are shown here. Image collected and cropped by CiteAb from the following publication

(https://journals.plos.org/plosone/article?

id=10.1371/journal.pone.0106155), licensed under a CC-BY license.

Immunohistochemistry-Paraffin: Smad7 Antibody [NBP2-24710] - IHC analysis of a formalin fixed paraffin-embedded (FFPE) human liver using Smad7 antibody on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images was performed using Aperio AT2 (Leica Biosystems). Cytoplasmic staining of Smad7 was observed. Staining was performed by Histowiz.

Western Blot: Smad7 Antibody [NBP2-24710] - Analysis using SMAD7 antibody. Lysate from HepG2 cells probed with SMAD7 antibody.

g		
g		
ry	1.1.4	
f	1 1	
	MW (kDa) 200 —	
	$^{116}_{97} \equiv$	
	66 — 55 —	
	36 —	SMAD7
	31 —	
	21	

Western Blot: Smad7 Antibody [NBP2-24710] - Analysis using the Azide Free version of NBP2-24710. Detection of lysate from A549 cells probed with SMAD7 antibody.



6

66 55

36 · 31 · 21 · 14 · Smad7

#### **Publications**

Takafumi Shichijo, Jun-ichirou Yasunaga, Kei Sato, Kisato Nosaka, Kosuke Toyoda, Miho Watanabe, Wenyi Zhang, Yoshio Koyanagi, Edward L. Murphy, Roberta L. Bruhn, Ki-Ryang Koh, Hirofumi Akari, Terumasa Ikeda, Reuben S. Harris, Patrick L. Green, Masao Matsuoka Vulnerability to APOBEC3G linked to the pathogenicity of deltaretroviruses Proceedings of the National Academy of Sciences of the United States of America 2024-03-19 [PMID: 38502701]

Tang Y, Du E, Wang G et al. A negative feedback loop centered on SMAD3 expression in transforming growth factor ?1-induced corneal myofibroblast differentiation Experimental eye research 2023-09-19 [PMID: 37734427] (WB, Human)

Shrestha R, Mohankumar K, Safe S Bis-indole derived nuclear receptor 4A1 (NR4A1) antagonists inhibit TGF betainduced invasion of embryonal rhabdomyosarcoma cells American journal of cancer research 2020-08-01 [PMID: 32905449] (WB, Human)

Yang P, Dong X, Zhang Y MicroRNA profiles in plasma samples from young metabolically healthy obese patients and miRNA-21 are associated with diastolic dysfunction via TGF- beta 1/Smad pathway J. Clin. Lab. Anal. 2020-02-28 [PMID: 32108968] (ELISA, Human)

Tang Y, Reissig S, Glasmacher E et al. Alternative Splice Forms of CYLD Mediate Ubiquitination of SMAD7 to Prevent TGFB Signaling and Promote Colitis Gastroenterology 2019-02-01 [PMID: 30315770] (WB, Mouse)

Kayalar O, Oztay F. Retinoic acid induced repair in the lung of adult hyperoxic mice, reducing transforming growth factor-B1 (TGF-B1) mediated abnormal alterations. Acta Histochem. [PMID: 24576683] (WB, Mouse)

Details:

Citation using the Azide Free form of this antibody.

Kleiter Ingo, Pedre Xiomara, Mueller Andre M et al. Inhibition of Smad7, a negative regulator of TGF-beta signaling, suppresses autoimmune encephalomyelitis. J Neuroimmunol. 2007-07-01 [PMID: 17553571]

Singha PK, Pandeswara S, Geng H et al. TGF-b induced TMEPAI/PMEPA1 inhibits canonical Smad signaling through R-Smad sequestration and promotes non-canonical PI3K/Akt signaling by reducing PTEN in triple negative breast cancer. Genes Cancer. 2014-09-01 [PMID: 25352949] (WB, Human)

Chang W, Wei K, Ho L et al. A Critical Role for the mTORC2 Pathway in Lung Fibrosis PLoS ONE 2014-08-28 [PMID: 25162417] (WB, Human)

Details:

SMAD7 antibody used for WB on lysates of fibroblasts (isolated from the surgical lung explants 0 Idiopathic Pulmonary Fibrosis /IPF) stimulated with TGF beta and treated or not with inhibitors MLN0128 (0.2 uM), PP242 (2 uM), or rapamycin (0.02 uM).

Roman-Blas JA, Stokes DG, Jimenez SA. Modulation of TGF-beta signaling by proinflammatory cytokines in articular chondrocytes. Osteoarthritis Cartilage. 2007-12-01 [PMID: 17604656]

L0nn P, Vanlandewijck M, Raja E et al. Transcriptional induction of salt-inducible kinase 1 by transforming growth factor B leads to negative regulation of type I receptor signaling in cooperation with the Smurf2 ubiquitin ligase. J Biol Chem. 2012-04-13 [PMID: 22378783]

Kidd M, Modlin IM, Pfragner R et al. Small bowel carcinoid (enterochromaffin cell) neoplasia exhibits transforming growth factor-beta1-mediated regulatory abnormalities including up-regulation of C-Myc and MTA1. Cancer. 2007-06-15 [PMID: 17469181]

Details:

WB: SMAD4 (IMG-565) and B-actin (IMG-531). Neoplastic kRJ-I enterochromaffin (EC) cells stimulated with growth factor-B1 (TGFB1), Fig 4.

More publications at <a href="http://www.novusbio.com/NBP2-24710">http://www.novusbio.com/NBP2-24710</a>





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-24710

NBP2-24710PEP	Smad7 Antibody Blocking Peptide
NBP2-24891	Rabbit IgG Isotype Control
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24710

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

