

Product Datasheet

TLR7 Antibody - BSA Free NBP2-24767

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-24767

Updated 2/23/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-24767




NBP2-24767

TLR7 Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein G purified
Buffer	PBS pH 7.2
Product Description	
Host	Rabbit
Gene ID	51284
Gene Symbol	TLR7
Species	Human, Primate, Rhesus Macaque
Reactivity Notes	Chimpanzee (100%), Dog (83%)
Immunogen	A portion of amino acids 900-950 of human TLR7 was used as the immunogen.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.5-2 ug/ml, Flow Cytometry 1-2 ug / 10 ⁶ cells, Immunohistochemistry, Immunohistochemistry-Paraffin reported in scientific literature (PMID 25586463)

Images


Flow Cytometry: TLR7 Antibody [NBP2-24767] - Intracellular staining by analysis of TLR7 in human PBMCs using this antibody at 2 ug/ml. Shaded histogram represents cells alone; green represents rabbit IgG isotype control, and red represents the TLR7 antibody.



Western Blot: TLR7 Antibody [NBP2-24767] - Analysis of human TLR7 in Ramos cell lysate in the 1) absence and 2) presence of immunizing peptide using this antibody.



Immunohistochemistry-Paraffin: TLR7 Antibody [NBP2-24767] - Human brain using 1:50 dilution of TLR7 antibody on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images (20X) was performed using Aperio AT2 (Leica Biosystems). Punctate staining of neurons was observed. Staining was performed by Histowiz.



Publications

Singh K, Agrawal NK, Gupta SK et al. Increased expression of endosomal members of toll-like receptor family abrogates wound healing in patients with type 2 diabetes mellitus *Int Wound J.* 2015-01-14 [PMID: 25586463] (WB, IHC-P, Human)

Details:

TLR7 antibody used for WB and IHC-P in study involving wound healing in patients with type 2 diabetes mellitus (T2DM)/ diabetic foot ulcer (DFU) patients in comparison to control wounds. IHC-P - Bouin's solution fixed paraffin embedded tissue sections from wound tissue samples, primary antibody used at 1:75 dilution in PBS with 4C 12 hours incubation in a wet chamber, detection with Vectastain Elite ABC Kit followed by hematoxylin counterstaining, data analysis included cells with brown-stained cytoplasm as positive for TLR7 (Figure 6). WB - whole tissue extracts of wound biopsies from DFU cases and controls, 50 ug protein ran on 10% gels, blocking with 5% of skimmed milk, primary antibody used at 1:1000 dilution, detection with HRP conjugated secondary followed by ECL development (Figure 7). These experiments showed increased TLR7 and TLR9 levels in wounds from T2DM patients compared with controls.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-24767

NBP2-26228-1mg	Imiquimod, TLR7 ligand
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24767

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

