

Product Datasheet

CD3 Antibody (OKT3) - BSA Free NBP2-25186

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 10

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-25186

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-25186



NBP2-25186

CD3 Antibody (OKT3) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	OKT3
Preservative	0.02% Sodium Azide
Isotype	IgG2a Kappa
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	23.1 kDa

Product Description	
Host	Mouse
Gene ID	916
Gene Symbol	CD3E
Species	Human, Mouse
Immunogen	This CD3 antibody was developed against human PBMCs.


Product Application Details	
Applications	Western Blot, Flow Cytometry, Functional, Immunoblotting, In vitro assay, In vivo assay
Recommended Dilutions	Western Blot reported in scientific literature (PMID 28699640), Flow Cytometry, Functional, Immunoblotting reported in scientific literature (PMID 23880762), In vitro assay, In vivo assay

Images


Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis using Alexa Fluor (R) 488 conjugate of NBP2-24867. A cell surface stain was performed on hPBMCs with CD8 Alpha antibody (RPA-T8) NBP2-25195 (top image) and a matched isotype control NBP2-27287 (bottom image). Cells were incubated in an antibody dilution of 1:200 for 20 minutes at room temperature. Both antibodies were conjugated to PerCP. A co-stain was also performed using CD3 antibody NBP2-24867AF488.




Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis using the Alexa Fluor(R) 700 conjugate of NBP2-24867. Staining of human PBMC. Flow cytometry image submitted by a verified customer review.




Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis using Azide Free version of NBP2-25186. CD3+ hPBMCs using the APC conjugate of this antibody. A cell surface stain was performed on CD3+ hPBMCs with V alpha 24 J alpha 18 TCR antibody (6B11) NBP2-00267PE (top image) and a matched isotype control




Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis of CD3 expression in hPBMC's gated to lymphocyte population at 1×10^6 cells/mL. Cells were stained with NBP2-24867PE (orange) conjugate at a 1:500 dilution. Results are shown with mIgG (PE conjugate) isotype control (blue). Image using the PE format of this antibody.



Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis of PerCP conjugate of NBP2-25186. A surface stain was performed on Jurkat cells with CD3 (OKT3) antibody NBP2-24867PCP (blue) and a matched isotype control (orange). Cells were incubated in an antibody dilution of 2.5 ug/mL for 20 minutes at room



Flow Cytometry: CD3 Antibody (OKT3) [NBP2-25186] - Analysis using the Alexa Fluor(R) 488 conjugate of NBP2-24867. Detection of CD3 expression in hPBMc's gated to lymphocyte population at 1×10^6 cells/mL. Cells were stained with NBP2-24867AF488 (orange) conjugate at a 1:500 dilution. Results are shown with mIgG2 (AF488 conjugate) isotype control (blue). Image using the Low Endotoxin and Azide Free form of this antibody.



Publications

Wu Q, Hu Y, Ma Q et al. Comprehensive Analysis of RAPGEF2 for Predicting Prognosis and Immunotherapy Response in Patients with Hepatocellular Carcinoma *Journal of Oncology* 2022-04-26 [PMID: 35518785] (FLOW)

Raab M, Lu Y, Kohler K et al. LFA-1 activates focal adhesion kinases FAK1/PYK2 to generate LAT-GRB2-SKAP1 complexes that terminate T-cell conjugate formation *Nat Commun* 2017-07-12 [PMID: 28699640] (WB, Human)

Ma SD, Xu X, Jones R et al. PD-1/CTLA-4 Blockade Inhibits Epstein-Barr Virus-Induced Lymphoma Growth in a Cord Blood Humanized-Mouse Model. *PLoS Pathog.* [PMID: 27186886] (In Vivo, Mouse)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Whitehurst CB, Li G, Montgomery SA et al. Knockout of Epstein-Barr Virus BPLF1 Retards B-Cell Transformation and Lymphoma Formation in Humanized Mice. *MBio.* [PMID: 26489865] (In Vivo, Human)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Kim SG, Lowe EL, Dixit D et al. Cocaine-mediated impact on HIV infection in humanized BLT mice. *Sci Rep.* [PMID: 26084721] (Func, In vitro, Human)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Ma SD Xu X Plowshay J et al. LMP1-deficient Epstein-Barr virus mutant requires T cells for lymphomagenesis. *J Clin Invest.* [PMID: 25485679] (In Vivo, Mouse)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Ma SD, Xu X, Plowshay J et al. LMP1-deficient Epstein-Barr virus mutant requires T cells for lymphomagenesis. *J Clin Invest.* [PMID: 25535359] (In Vivo)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Gill T, Levine AD. Mitochondria-derived hydrogen peroxide selectively enhances T cell receptor-initiated signal transduction. *J Biol Chem.* [PMID: 23880762]

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Crellin NK Garcia RV Hadisfar O et al. Human CD4+ T cells express TLR5 and its ligand flagellin enhances the suppressive capacity and expression of FOXP3 in CD4+CD25+ T regulatory cells. *J Immunol.* [PMID: 16339542] (FLOW, Human)

Details:

Citation using the Azide and Endotoxin Free form of this antibody.

Ma SD, Yu X, Mertz JE et al. An Epstein-Barr Virus (EBV) mutant with enhanced BZLF1 expression causes lymphomas with abortive lytic EBV infection in a humanized mouse model. *J Virol.* 2012-08-01 [PMID: 22623780] (Mouse)

Details:

CD3 (IMG-6240E). FA: Mice received 2 ug of CD3 antibody intravenously per day, starting 1 day after EBV infection, Fig 1A & Fig 1B.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-25186

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-96981-0.5mg	Mouse IgG2a Kappa Isotype Control (M2AK)
NBP2-24867G	CD3 Antibody (OKT3) [DyLight 488]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-25186

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

