

Product Datasheet

IL-17/IL-17A Antibody (4H1524) - Azide Free NBP2-25229

Unit Size: 0.25 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-25229

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-25229



NBP2-25229

IL-17/IL-17A Antibody (4H1524) - Azide Free

Product Information	
Unit Size	0.25 mg
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4H1524
Preservative	No Preservative
Isotype	IgG2b Kappa
Purity	Protein G purified
Buffer	0.2 ml sterile PBS


Product Description	
Host	Mouse
Gene ID	3605
Gene Symbol	IL17A
Species	Human
Immunogen	A portion of amino acids 1-100 was used as immunogen.

Product Application Details	
Applications	Western Blot, Flow Cytometry, Block/Neutralize, CyTOF-ready
Recommended Dilutions	Western Blot 1:100-1:2000, Flow Cytometry 1ul/1 million cells, CyTOF-ready, Block/Neutralize




Images


Western Blot: IL-17/IL-17A Antibody (4H1524) - Azide Free [NBP2-25229] - Analysis of IL-17A in (1) human, (2) mouse and (3) rat full-length recombinant IL-17A protein using NBP2-25229 at 1 ug/ml.



Flow Cytometry: IL-17/IL-17A Antibody (4H1524) - Azide Free [NBP2-25229] - Analysis using the PE conjugate of NBP2-25229. IL-17A positive cells in PMA/IM stimulated fresh PBMC: PBMC stimulated for 6h with PMA (25ng/ml) & IM (250ng/ml), in the presence of BFA. Cells were surface stained with CD4 APC, fixed & permeabilized for intracellular staining with PE conjugated



Block/Neutralize: IL-17/IL-17A Antibody (4H1524) - Azide Free [NBP2-25229] - Neutralization by Human IL-17A Antibody. Secretion of IL-6 by 3T3 cells stimulated with recombinant human IL-17A (20ng/ml) is neutralized by increasing concentrations of IL-17A monoclonal antibody. Antibody at different concentrations was added to cells 1hr prior to addition of cytokine. Supernatant collected from 60hr cultures was used for analysis to measure secreted IL-6 by ELISA.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-25229

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43317-0.5mg	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b)
NBP2-27252	IL-17/IL-17A Antibody (4H1524) [PE]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-25229

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

