

Product Datasheet

RelA/NFkB p65 Antibody (112A1021) - Azide and BSA Free NBP2-27416

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-27416

Updated 4/13/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-27416



NBP2-27416

RelA/NFkB p65 Antibody (112A1021) - Azide and BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	112A1021
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	0.2 ml sterile PBS

Product Description	
Host	Mouse
Gene ID	5970
Gene Symbol	RELA
Species	Human, Mouse, Rat
Reactivity Notes	Predicted to react with Bovine, Gorilla, New World Monkey and Porcine.
Immunogen	This Nfkb p65 antibody was raised against a synthetic peptide corresponding to amino acids 526-539 of human NF-kB (p65) protein.


Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot 2-5ug/ml~. Use reported in scientific literature (Wittmann et al (2005)), Simple Western, Flow Cytometry 1ul/1 million cells, Immunohistochemistry, Immunohistochemistry-Paraffin 5ug/ml. Use reported in scientific literature (Chang et al (2005)), Flow (Intracellular) reported in scientific literature (PMID 15955098), CyTOF-ready
Application Notes	In Simple Western only 10-15 uL of the recommended dilution is used per data point.

Images


Western Blot: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - Lysates of HeLa human cervical epithelial carcinoma cell line and Daudi human Burkitt's lymphoma cell line. PVDF membrane was probed with 1 ug/mL mouse anti-RelA/NFkB p65 monoclonal (NB100-56712, Novus Biologicals), followed by 1:2000 dilution of the appropriate HRP-conjugated secondary antibody, donkey anti-mouse IgG (HAF018). Image using the standard format of this product.




Immunohistochemistry-Paraffin: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - Formalin-fixed, paraffin-embedded ovarian cystadenocarcinoma probed with p65 antibody at 5 ug/ml. Human tissue TMA was used for this test. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.




Flow Cytometry: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - Intracellular staining of 293 HEK cells using 0.5 ug of p65 antibody. Green histogram represents the isotype control (p65) antibody. this antibody was used for this test, and an anti-mouse IgG PE conjugated secondary antibody.




Western Blot: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - analysis of p65 using p65 antibody at 2 ug/ml in 30 ug of A) Ramos, B) Daudi, C) HeLa and D) mouse NIH 3T3 cell lysate.



Simple Western: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - Simple Western lane view shows a specific band for NFkB p65 in 1.0 mg/ml of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Simple Western: RelA/NFkB p65 Antibody (112A1021) - Azide Free [NBP2-27416] - Detection of p65 by Simple Western (JESS) in chemiluminescence in nuclear extract of reconstructed human epidermis lysate (about 600 ug/mL protein concentration). Antibody at 1:20. Simple Western image submitted by a verified customer review. Image using the standard format of this product.



Publications

Wittmann M, Purwar R, Hartmann C et al. Human keratinocytes respond to interleukin-18: implication for the course of chronic inflammatory skin diseases. *J Invest Dermatol.* 2005-06-01 [PMID: 15955098] (Flow Cytometry Control, Human)

Details:

Flow (Intracellular), Fig 3 (Intact nuclei isolated from normal human keratinocytes: analysis of NF-kB translocation to the nucleus).

Santhi WS, Sebastian P, Varghese BT et al. NF-kappaB and COX-2 during oral tumorigenesis and in assessment of minimal residual disease in surgical margins. *Exp Mol Pathol.* 2006-10-01 [PMID: 16822500] (IHC-P, Human)

Details:

Products used (human): 1. IkbA [IMG-127A]: IHC paraffin (normal oral mucosa, leukoplakia, and oral cancer), Fig. 1 IHC paraffin analysis (normal oral mucosa, leukoplakia, and oral cancer), Fig. 2 IHC paraffin (oral cancer: primary tumor, surgical margin,

Chang MS, Lee HS, Jung EJ et al. Cell-cycle regulators, bcl-2 and NF-kappaB in Epstein-Barr virus-positive gastric carcinomas. *Int J Oncol.* 2005-11-01 [PMID: 16211221] (IHC-P, Human)

Details:

NF-kB (p65), IMG-150A: IHC (paraffin), Fig 2F (human gastric carcinoma).

Lee TL, Yeh J, Friedman J et al. A signal network involving coactivated NF-kappaB and STAT3 and altered p53 modulates BAX/BCL-XL expression and promotes cell survival of head and neck squamous cell carcinomas. *Int J Cancer.* 2008-05-01 [PMID: 18172861]

Details:

IMGEX products cited: 1. NF-kB p65 mAb-biotinylated, clone 112A1021 (IMG-150B): IP (UM-SCC human squamous carcinoma cell lines), Fig 2b. Immunoprecipitates were western blotted with the purified format of the p65 mAb clone 112A1021 (IMG-150A) or with a p



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-27416

NBL1-13619	NFkB p105/p50 Overexpression Lysate
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-27416

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

