Product Datasheet

TIMP-3 Antibody NBP2-29478

Unit Size: 0.1 mg

Store at 4C.



Reviews: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-29478

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-29478



NBP2-29478


TIMP-3 Antibody

Product Information Unit Size 0.1 mg Concentration 0.2 mg/ml Storage Store at 4C. Clonality Polyclonal Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 30 kDa Product Description 200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to a-80C. Host Rabbit Gene ID 7078 Gene Symbol TiMP3 Species Human, Mouse, Rat, Bovine, Canine, Equine Reactivity Notes The amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review. Immunosen TIMP-3 (aa175-211) (Uniprot: P35625) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofistochemistry 1-2ug/ml, Immunohistochemistry - Paraffin Recommended Dilutions Flow Cytometry. Immun			
Concentration0.2 mg/mlStorageStore at 4C.ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityProtein A purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is nowse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2 ug/mil, Immunohistochemistry/ Immunohistochemistry 1-2 ug/mil, Immunohistochemistry 1-2 ug/mil Immunohistochemistry 1-2 ug/mil, Immunohistochemistry Immunohistochemistry 1-2 ug/mil, Immunohistochemistry Immunohistochemistry 1-2 ug/mil, Immunohistochemistry 1-2 ug/mil Immunohistochemistry 1-2 ug/mil, Immunohistochemistry 1-2 ug/mil	Product Information		
StorageStore at 4C.ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityProtein A purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is nouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunofluorescence, Immunofluorescence, Immunohistochemistry, Immunofluorescence, 1-2 ug/ml, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry-ParaffinApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with TMM	Unit Size	0.1 mg	
ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityProtein A purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mg/ml. (NBP2-33098)Poscription200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mg/ml. (NBP2-33098)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunofistochemistry-Immunofistochemistry-ParaffinRecommended DilutionsFlow Cytometry, Immunofistochemistry-ParaffinRecommended DilutionsFlow Cytometry, Immunofistochemistry - 2ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heading tissues sections in 10mM Tris with 1mMEDTA, p.H. 0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	0.2 mg/ml	
Preservative 0.05% Sodium Azide Isotype IgG Purity Protein A purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 30 kDa Product Description 200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7078 Gene Symbol TiMP3 Species Human, Mouse, Rat, Bovine, Canine, Equine Reactivity Notes TiMP3 (an175-211) (Uniprot: P35625) Product Application Details Flow Cytometry 1-2 ug/mil mnunocytochemistry/ Immunofluorescence, Immunohistochemistry -Paraffin Recommended Dilutions Flow Cytometry 1-2 ug/mil mnunofluorescence 1-2 ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunocytochemistry/ Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Storage	Store at 4C.	
IsotypeIgGPurityProtein A purifiedBuffer10 mM PBS with 0.05% BSATarget Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098)Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofistochemistry 1-2ug/ml, Immunocytochemistry, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/ Immunofistochemistry 1-2ug/ml, Immunohistochemistry I aug/ml, Immunohistochemistry 1-2ug/ml, Immunohistochemistry I-2ug/ml, Immunohistochemistry I-2ug/m	Clonality	Polyclonal	
Purity Protein A purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 30 kDa Product Description 200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7078 Gene Symbol TIMP3 Species Human, Mouse, Rat, Bovine, Canine, Equine Reactivity Notes The amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review. Immunogen TIMP-3 (aa175-211) (Uniprot: P35625) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Flow Cytometry 1-2 ug/million cells, Immunohistochemistry-Paraffin Recommended Dilutions Flow Cytometry (Formalin-fixed): 1-2ug/ml of 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10M Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide	
Buffer10 mM PBS with 0.05% BSATarget Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunofluorescence 1-2 ug/ml, Immunohistochemistry I-2ug/ml, Immunohistochemistry I-2ug/ml, Immunohistochemistry I-2ug/ml, Immunohistochemistry-Paraffin Paraffin 1-2 ug/mlApplication NotesFlow Cytometry 1-2 ug/mlilion cells, Immunohistochemistry-Paraffin EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG	
Target Molecular Weight30 kDaProduct Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunofstochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry -ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry -Paraffin 1-2 ug/mlApplication NotesFlow Cytometry 1-2 ug/million cells, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesFlow Cytometry 1-2 ug/million cells, Immunohistochemistry - Paraffin 1-2 ug/ml	Purity	Protein A purified	
Product Description Description 200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7078 Gene Symbol TIMP3 Species Human, Mouse, Rat, Bovine, Canine, Equine Reactivity Notes The amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review. Immunogen TIMP-3 (aa175-211) (Uniprot: P35625) Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryParaffin Recommended Dilutions Flow Cytometry 1-2 ug/million cells, Immunohistochemistry -Paraffin Recommended Dilutions Flow Cytometry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed issues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA	
Description200ug/ml of antibody purified from rabbit anti-serum by Protein A. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry -ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunohistochemistry (Formalin-fixed): 1-2ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	30 kDa	
MM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry Paraffin 1-2 ug/milApplication NotesFlow Cytometry formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining go formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description		
HostRabbitGene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0 mg/ml. (NBP2-33098) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to	
Gene ID7078Gene SymbolTIMP3SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, 	Host		
SpeciesHuman, Mouse, Rat, Bovine, Canine, EquineReactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	7078	
Reactivity NotesThe amino acid sequence used as the immunogen for the TIMP3 antibody is 100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	TIMP3	
100% homologous in human, cow, dog and horse, and 94% homologous in mouse and rat. Sheep data from customer review.ImmunogenTIMP-3 (aa175-211) (Uniprot: P35625)Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human, Mouse, Rat, Bovine, Canine, Equine	
Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Reactivity Notes	100% homologous in human, cow, dog and horse, and 94% homologous in	
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	TIMP-3 (aa175-211) (Uniprot: P35625)	
Immunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsFlow Cytometry 1-2 ug/million cells, Immunohistochemistry 1-2ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details		
Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry- Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications		
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-	
	Application Notes	of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	



Images

Immunohistochemistry-Paraffin: TIMP-3 Antibody [NBP2-29478] - TIMP-3 is expressed in first trimester human decidual cells. This image was submitted via customer Review.



Immunohistochemistry-Paraffin: TIMP-3 Antibody [NBP2-29478] -Formalin-fixedn, paraffin-embedded human placenta stained with TIMP3 Rabbit polyclonal antibody





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-29478

210-TA-005	TNF-alpha [Unconjugated]
NBP2-24891	Rabbit IgG Isotype Control
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-29478

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

