Product Datasheet

NF-H Antibody (RT97) - Azide and BSA Free NBP2-33062

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-33062

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-33062



NBP2-33062


NF-H Antibody (RT97) - Azide and BSA Free

WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy sunit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H) Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	NF-H Antibody (RT97) - Azide and BSA Free		
Concentration 1.0 mg/ml	Product Information		
Storage Store at -20 to -80C. Avoid freeze-thaw cycles.	Unit Size	0.1 mg	
Clone RT97 Preservative No Preservative Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 200 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-M) and sympathetic ganglion cells. Neurofilaments one in eural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilaments. Neuroflaments as also present in paragangliomas as well as adfenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Concentration	1.0 mg/ml	
Clone	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.	
Preservative IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 200 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H) Anti-neurofilament stains a number of neural, neuroendocrine, and endocriot tumors. Neuromas, ganglioneuromas, gangliongemas, agnipioneurofilament. Neurofilament stains a number of neural, neuroendocrine, and endocriot tumors. Neuromas, ganglioneuromas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Clonality	Monoclonal	
IgG1 Kappa	Clone	RT97	
Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 200 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10 mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major suburnithm rolecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-M) Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogimas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilaments. Neurofilaments also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilaments. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Preservative	No Preservative	
Buffer 10 mM PBS Target Molecular Weight 200 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and at cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Isotype	IgG1 Kappa	
Target Molecular Weight 200 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy sunit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, ganglioneuromas, anglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Purity	Protein A or G purified	
Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H) Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, ganglioneurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Buffer	10 mM PBS	
1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H) Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Target Molecular Weight	200 kDa	
Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/m WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 4744 Gene Symbol NEFH Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Product Description		
Gene Symbol Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Description	Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-29435). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to	
Gene Symbol Species Human, Mouse, Rat, Porcine, Chicken Marker Neuronal Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Host	Mouse	
Species	Gene ID	4744	
Marker Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliojiomas, ganglioneuroblastoma and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Gene Symbol	NEFH	
Specificity/Sensitivity This monoclonal antibody reacts with a 200kDa protein, identified as heavy sunit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Species	Human, Mouse, Rat, Porcine, Chicken	
unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerve and sympathetic ganglion cells. Neurofilaments consist of three major subur with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H Anti-neurofilament stains a number of neural, neuroendocrine, and endocrin tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastom and neuroblastomas stain positively for anti-neurofilament. Neurofilaments a also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, at oat cell carcinomas of the lung also express neurofilament. Immunogen Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036) Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Marker	Neuronal Marker	
Product Application Details Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin		elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.	
Applications Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Immunogen	Triton-X 100 insoluble protein fraction of rat brain (Uniprot: P12036)	
Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	Product Application Details		
	Applications	Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	
Application Notes Immunohistochemistry (Formalin-fixed): 0.25-0.5ug/ml for 30 minutes at RT.	Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin	
Staining of formalin-fixed tissues requires heating tissue sections in 10mM T with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.	Application Notes	minutes.	




Images


Western Blot: NF-H Antibody (RT97) - Azide and BSA Free [NBP2-33062] - Western Blot Analysis of human brain tissue lysate using NF-H Antibody (RT97).



Immunohistochemistry-Paraffin: NF-H Antibody (RT97) - Azide and BSA Free [NBP2-33062] - Formalin-fixed, paraffin-embedded Rat Cerebellum stained with NF-H Antibody (RT97).



Immunohistochemistry-Paraffin: NF-H Antibody (RT97) - Azide and BSA Free [NBP2-33062] - Formalin-fixed, paraffin-embedded human Cerebellum stained with NF-H Antibody (RT97).



Publications

Reincke SM, von Wardenburg N, Homeyer MA et al. Chimeric autoantibody receptor T cells deplete NMDA receptor-specific B cells Cell 2023-10-26 [PMID: 37918394] (IHC-P, Mouse)

Details:

1:200 dilution





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

rax. 905.027.0402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-33062

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
H00004744-Q01-10ug Recombinant Human NF-H GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-33062

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

