Product Datasheet

CD45RA Antibody (158-4D3) [Allophycocyanin/Cy7] NBP2-33144APCCY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-33144APCCY7

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP2-33144APCCY7



NBP2-33144APCCY7

T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RO and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation	CD45RA Antibody (158-4D3) [Allophycocyanin/Cy7]		
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information		
Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone 158-4D3 Preservative 0.05% Sodium Azide Isotype IgG2a Kappa Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 5788 Gene ID 5788 Gene Symbol PTPRC Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.8], CD45RA is isoforms of the human leukocyte common antigen (CD45), Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA are naive or virgin T-cells. To-cells expressing CD45RA are naive or virgin T-cells. To-cell expressing CD45RA are naive or virgin T-cells. To-cell expressing CD45RA are naive or virgin T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. Sin monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. Sin monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. In the subpopulation of CD4+ or CD8+ T-cells. The monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. The monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. The monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. The monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antibody is useful to the subpopulation of CD4+ or CD8+ T-cells. This monoclonal antib	Unit Size	0.1 ml	
Clonality Monoclonal Clone 158-4D3 Preservative 0.05% Sodium Azide Isotype IgG2a Kappa Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 5788 Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.3B], CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD8+17-cells, 50% of peripheral CD8+10%. The coll expressing CD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing CD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing cD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing cD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing cD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing cD45RA are naive or virgin 17-cells, 50% of peripheral CD8+10%. The coll expressing cD45RA a	Concentration	·	
Preservative	Storage	Store at 4C in the dark. Do not freeze.	
Preservative 0.05% Sodium Azide	Clonality	Monoclonal	
IgG2a Kappa IgG2a Kappa Allophycocyanin/Cy7	Clone	158-4D3	
Conjugate Allophycocyanin/Cy7 Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38], CD45RA is isoforms of the human leukocyte common antigen (CD45), Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed not-50% of peripheral CD4+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA ave). Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatior and permeabilization steps in your flow protocol may be required. Place contact our technical support team at technical@enovusbio.com if you have any	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO+ve) from B cell lymphomas (CD45RA+ve). Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunositochemistry, Immunocytochemistry/ or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Place contact our technical support team at technical@novusbio.com if you have any	Isotype	IgG2a Kappa	
Product Description	Conjugate	Allophycocyanin/Cy7	
Product Description Host Mouse Gene ID 5788 Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed n40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA are naive or virgin T-cells. T-cells captesing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. It is monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Purity	Protein A or G purified	
Mouse	Buffer	PBS	
Gene Symbol PTPRC Species Human Marker Leucocyte Marker Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA or a memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Product Description		
Species	Host	Mouse	
Species	Gene ID	5788	
Marker Leucocyte Marker Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells. B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA are naive or virgin T-cells. T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunostochemistry, Immunocytochemistry/ Immunocytochemistry/ Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatior and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Gene Symbol	PTPRC	
Specificity/Sensitivity Recognizes a protein of 205kDa-220kDa, identified as CD45RA [Workshop V; Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RA are naive or virgin T-cells. T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry/ Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Recommended Dilutions Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatior and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Species	Human	
Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve). Immunogen Stimulated human leukocytes Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixatior and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Marker	Leucocyte Marker	
Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Specificity/Sensitivity	Code CD45.38]. CD45RA is isoforms of the human leukocyte common antigen (CD45). Human CD45 contains three exons which encode peptide segments designated A, B and C, respectively. The differential splicing of the exons generates at least five isoforms, ABC, AB, BC, B and O. This antibody reacts with ABC and BC isoforms. CD45RA is expressed on 40-50% of peripheral CD4+ T-cells, 50% of peripheral CD8+ T-cells, B-cells, and leukemic B-cell lines. T-cells expressing CD45RA are naive or virgin T-cells. T-cells expressing CD45RO are memory T-cells. CD45RA and CD45RO define complementary, predominantly non-overlapping populations of resting peripheral T-cells. This monoclonal antibody is useful in study on the subpopulation of CD4+ or CD8+ T-cells. It can especially be used to differentiate T-cell lymphomas (CD45RO +ve) from B cell lymphomas (CD45RA +ve).	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Immunogen	Stimulated human leukocytes	
Immunohistochemistry, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any			
Application Notes Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Applications		
optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	Recommended Dilutions		
	Application Notes	optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-33144APCCY7

NBP1-96981APCCY7 Mouse IgG2a Kappa Isotype Control (M2AK) [Allophycocyanin/Cy7]

210-TA-005 TNF-alpha [Unconjugated]

M6000B-1 IL-6 [HRP]

6507-IL-010/CF IL-4 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-33144APCCY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

