

Product Datasheet

CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free NBP2-33154

Unit Size: 0.1 mg

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-33154

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-33154



NBP2-33154

CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	C31.3
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS
Target Molecular Weight	82.5 kDa

Product Description	
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-15202). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	5175
Gene Symbol	PECAM1
Species	Human
Reactivity Notes	Weakly reacts with Rat.
Marker	Endothelial Cell Marker
Specificity/Sensitivity	CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.
Immunogen	Recombinant full-length human CD31/PECAM-1 protein (Uniprot: P16284)

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, CyTOF-ready
Recommended Dilutions	Western Blot 0.5-1.0ug/ml, Flow Cytometry 0.5-1ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 0.5-1ug/ml, Immunohistochemistry-Paraffin 0.5-1.0ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml, CyTOF-ready




Application Notes


Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images


Western Blot: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Western Blot Analysis of human Jurkat cell lysate using CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free.




Immunocytochemistry/Immunofluorescence: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Flow Cytometry of human CD31 on Jurkat cells. Black: cells alone; Grey: Isotype Control; Green: AF488-labeled CD31/PECAM-1 Antibody (C31.7) - Azide and BSA Free.




Immunohistochemistry-Paraffin: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Formalin-paraffin human Angiosarcoma stained with CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free.




Flow Cytometry: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Flow Cytometric analysis of human CD31 on Jurkat cells. Black: cells alone; grey: Isotype Control; Green: AF488-labeled CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free.



Immunohistochemistry-Paraffin: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Staining with CD31/PECAM-1 Antibody (C31.3) [Alexa Fluor® 647] (NBP2-33154AF647) using GE Cell DIVE (20x) on Human Rheumatoid Arthritis synovium (10 ug/ml). Image from verified customer review.



Immunohistochemistry-Paraffin: CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free [NBP2-33154] - Formalin-paraffin human tonsil stained with CD31/PECAM-1 Antibody (C31.3) - Azide and BSA Free.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-33154

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-54655PEP	CD31/PECAM-1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-33154

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

