

# Product Datasheet

## p107 Antibody - BSA Free NBP2-33735

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-33735](http://www.novusbio.com/NBP2-33735)

Updated 2/23/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-33735](http://www.novusbio.com/reviews/destination/NBP2-33735)



**NBP2-33735**

p107 Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol

Product Description	
Host	Rabbit
Gene ID	5933
Gene Symbol	RBL1
Species	Human
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non-tested species: Mouse (86%), Rat (86%)
Immunogen	This antibody was developed against a recombinant protein corresponding to amino acids: ESLAWSHDSALWEALQVSANKVPTCEEVIFPNNFETGNGGNVQGHPLPLMPMS PLMHPRVKEVRTDSGSLRRDMQPLSPISVHER


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:200 - 1:500, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:200 - 1:500
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.

**Images**


Immunohistochemistry-Paraffin: p107 Antibody [NBP2-33735] - Staining of human colon, kidney, liver and testis using Anti-p107 antibody NBP2-33735 (A) shows similar protein distribution across tissues to independent antibody NBP2-33791 (B).




Western Blot: p107 Antibody [NBP2-33735] - Lane 1: Marker [kDa] 250, 130, 95, 72, 55, 36, 28, 17, 10. Lane 2: Human cell line RT-4. Lane 3: Human cell line U-251MG sp




Immunocytochemistry/Immunofluorescence: p107 Antibody [NBP2-33735] - Selinexor alters pocket protein expression in nuclear and cytoplasmic compartments. Representative images of p107, p130 and RB (green) IF staining of bladder cancer cells treated with vehicle (V) or selinexor (S) for 48 hours. Tubulin staining (red) and DAPI staining (blue) served to define the cytoplasmic and nuclear compartments, respectively. The inserts are magnifications of the boxed cells. Image collected and cropped by CiteAb from the following publication (<https://www.oncotarget.com/fulltext/26179>), licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: p107 Antibody [NBP2-33735] - Immunofluorescent staining of human cell line HeLa shows localization to nucleoplasm.




Immunohistochemistry-Paraffin: p107 Antibody [NBP2-33735] - Staining of human testis shows moderate nuclear positivity in cells in seminiferous ducts and Leydig cells.




Immunohistochemistry-Paraffin: p107 Antibody [NBP2-33735] - Staining of human colon shows moderate nuclear positivity in glanular cells.




Immunohistochemistry-Paraffin: p107 Antibody [NBP2-33735] - Staining of human kidney shows moderate nuclear postivity in cells in tubules.




Immunohistochemistry-Paraffin: p107 Antibody [NBP2-33735] - Staining of human liver shows no positivity in hepatocytes as expected.



**Western Blot: p107 Antibody [NBP2-33735] - Selinexor alters pocket protein expression in nuclear & cytoplasmic compartments(A)** Representative images of p107, p130 & RB (green) IF staining of bladder cancer cells treated with vehicle (V) or selinexor (S) for 48 hours. Tubulin staining (red) & DAPI staining (blue) served to define the cytoplasmic & nuclear compartments, respectively. The inserts are magnifications of the boxed cells. (B) Quantification of staining intensity of pocket proteins normalized to DAPI. (C) Nuclear & cytoplasmic fractions of cell treated with vehicle or 0.15  $\mu$ M selinexor (UM-UC-3 & T24 cells), 0.25  $\mu$ M selinexor (J82) & 0.5  $\mu$ M selinexor (TCCSUP) for 72 hours were assessed for the expression of RB, p107 & p130. Nup62 & tubulin were used as markers for the nuclear & cytoplasmic fractions, respectively. (D) T24 & UM-UC-3 cells transfected with siC or siRB & were treated with vehicle or 0.1  $\mu$ M selinexor for 72 hours. The results are shown as percent cell viability comparing drug treated to vehicle treated cells. (E) Palbociclib reduces T24 & UM-UC-3 bladder tumor cells viability in a dose dependent manner. (F) Combined selinexor (0.1  $\mu$ M) & palbociclib (0.5  $\mu$ M) treatment is more effective in reducing viability of cells than either treatment alone where the CI = 1.04 for UM-UC-3 cells & 1.02 for T24 cells indicating an additive response. Error bars =  $\pm$  standard deviation. Student's t test; \* denotes  $p \leq 0.05$ , \*\* denotes  $p \leq 0.01$ . Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30349650>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



**Immunocytochemistry/ Immunofluorescence: p107 Antibody [NBP2-33735] - Selinexor alters pocket protein expression in nuclear & cytoplasmic compartments(A)** Representative images of p107, p130 & RB (green) IF staining of bladder cancer cells treated with vehicle (V) or selinexor (S) for 48 hours. Tubulin staining (red) & DAPI staining (blue) served to define the cytoplasmic & nuclear compartments, respectively. The inserts are magnifications of the boxed cells. (B) Quantification of staining intensity of pocket proteins normalized to DAPI. (C) Nuclear & cytoplasmic fractions of cell treated with vehicle or 0.15  $\mu$ M selinexor (UM-UC-3 & T24 cells), 0.25  $\mu$ M selinexor (J82) & 0.5  $\mu$ M selinexor (TCCSUP) for 72 hours were assessed for the expression of RB, p107 & p130. Nup62 & tubulin were used as markers for the nuclear & cytoplasmic fractions, respectively. (D) T24 & UM-UC-3 cells transfected with siC or siRB & were treated with vehicle or 0.1  $\mu$ M selinexor for 72 hours. The results are shown as percent cell viability comparing drug treated to vehicle treated cells. (E) Palbociclib reduces T24 & UM-UC-3 bladder tumor cells viability in a dose dependent manner. (F) Combined selinexor (0.1  $\mu$ M) & palbociclib (0.5  $\mu$ M) treatment is more effective in reducing viability of cells than either treatment alone where the CI = 1.04 for UM-UC-3 cells & 1.02 for T24 cells indicating an additive response. Error bars =  $\pm$  standard deviation. Student's t test; \* denotes  $p \leq 0.05$ , \*\* denotes  $p \leq 0.01$ . Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30349650>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Baek HB, Lombard AP, Libertini SJ et al. XPO1 inhibition by selinexor induces potent cytotoxicity against high grade bladder malignancies. *Oncotarget*. 2018-10-02 [PMID: 30349650] (WB, ICC/IF, Human)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-33735**

---

NBP2-33735PEP	p107 Recombinant Protein Antigen
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-33735](http://www.novusbio.com/reviews/submit/NBP2-33735)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

